arXiv:1604.01626v1 [physics.chem-ph] 6 Apr 2016

Room temperature chiral discrimination in paramagnetic NMR spectroscopy

Alessandro Soncini* and Simone Calvello
School of Chemistry, University of Melbourne, VIC 3010, Australia

A recently proposed theory of chiral discrimination in NMR spectroscopy based on the detection of a molecular electric polarization $\boldsymbol{P}$ rotating in a plane perpendicular to the NMR magnetic field [A. D. Buckingham, J. Chem. Phys. **140**, 011103 (2014)], is here generalized to paramagnetic systems. Our theory predicts new contributions to $\boldsymbol{P}$, varying as the square of the inverse temperature. Ab initio calculations for ten Dy^{3+} complexes, at 293K, show that in strongly anisotropic paramagnetic molecules $\boldsymbol{P}$ can be more than 1000 times larger than in diamagnetic molecules, making *paramagnetic* NMR chiral discrimination amenable to room temperature detection.

Despite its central role in biological processes and in a wide range of chemical reactions, chirality, the property of molecules lacking improper symmetry elements to be distinguishable from their mirror image (or enantiomer), remains a challenging property to detect and quantify [1], making the development of new spectroscopic techniques to achieve this goal a strategic research field [2–9].

Magnetic resonance spectroscopies, despite being among the most useful characterization techniques due to their high sensitivity to tiny details of the geometrical and electronic structure of molecules, are blind to chirality. However, it has been recently proposed by Buckingham [3, 9] and Fischer [5] that NMR could be used to achieve chiral discrimination for closed-shell chiral molecules via a minor modification of the experimental set up, so to make it fit for the detection of a rotating average electric polarization $\overline{\boldsymbol{P}}$ induced by the combined effect of the NMR magnetic field $\boldsymbol{B}$, and the nuclear magnetic dipole moment $\boldsymbol{m}^I$ associated to a nucleus I of the chiral molecule, rotating in the plane perpendicular to $\boldsymbol{B}$ following a resonant $\pi/2$ radiofrequency pulse. In particular, the induced $\overline{\boldsymbol{P}}$ is always oriented along $\boldsymbol{B}\times\boldsymbol{m}^I$ (thus rotating with $\boldsymbol{m}^I$), points in opposite directions for the two enantiomers (hence its chirality-sensitivity), and is proportional to the pseudoscalar $\sigma^{(1)}$, isotropic average of a third-rank tensor σ_{ijk} $(i,j,k=x,y,z)$ known as shielding polarizability [3, 5, 10]. However, computational estimates of $\sigma^{(1)}$ in diamagnetic molecules suggest that it is generally too small to be detected [9, 11, 12].

In this Letter we propose a theory of paramagnetic NMR chiral discrimination which is valid for molecules with a ground state of arbitrary degeneracy. We describe the response of the degenerate system in terms of a generalized shielding polarizability tensor Φ_{ijk} defined by analytical third–derivatives of the free energy, reducing to σ_{ijk} in the limit of a non-degenerate ground state. The proposed free-energy response theory features previously unexplored terms proportional to the square of the inverse temperature $\beta=1/k_BT$, so that $\sigma^{(1)}$ becomes $\sigma^{(1)}+\beta^2\sigma^{(1p)}$. Finally, we present ab initio calculations showing that $\beta^2\sigma^{(1p)}$ yields a contribution to $\overline{\boldsymbol{P}}$ at 293K that is orders of magnitude larger than in closed-shell molecules, potentially observable at room temperature.

Theory. Let us consider an ensemble of nuclear magnetic moments $\boldsymbol{m}^I=\mu_N g_I\boldsymbol{I}$, with μ_N the nuclear magneton, g_I the g-factor of the probed nucleus I, and $\boldsymbol{I}$ the nuclear spin in units of $\hbar$. In a pulsed NMR experiment, the nuclear moments $\boldsymbol{m}^I$, partially aligned along a magnetic field $\boldsymbol{B}=B_0\boldsymbol{u}_z$, are flipped in the xy-plane perpendicular to $\boldsymbol{B}$ by a radiofrequency $\pi/2$-pulse in resonance with the nuclear Larmor frequency $\omega^I=\mu_N g_I(1-\sigma)|\boldsymbol{B}|$. Here σ is the nuclear shielding constant, the isotropic average of the shielding tensor $\boldsymbol{\sigma}$ describing the departure from $\boldsymbol{B}$ of the local magnetic field at nucleus I due to surrounding electrons. The coherent in-plane precession of $\boldsymbol{m}^I$ gives rise to a free-induction decay (FID) signal picked up by a coil with axis in the xy-plane, whose Fourier transform provides the NMR spectrum.

The combined effect of $\boldsymbol{B}$ and $\boldsymbol{m}^I$ on the electrons of a diamagnetic molecule has been shown to induce an electric dipole polarization [3, 5, 9], which can be obtained by expanding the energy of the molecule $W(\boldsymbol{E},\boldsymbol{B},\boldsymbol{m}^I)\equiv W$ in a power series of $\boldsymbol{B}$, $\boldsymbol{m}^I$, and an external electric field $\boldsymbol{E}$, thermally averaged over all molecular orientations in the presence of $\boldsymbol{B}$ and $\boldsymbol{m}^I$ [9]. Truncation of the ensuing power series to leading order in the inducing fields yields the average induced polarization $\overline{P_i}\equiv\overline{P}_{d,i}$ as [3, 5, 9]:

$$\overline{P}_{d,i}=-\left\langle\frac{\partial W}{\partial E_i}\right\rangle\approx-\left\langle\sigma_{ijk}+\beta\bar{\mu}_i\sigma_{jk}\right\rangle_{\rm rot}B_j m_k^I \qquad (1)$$

where subscript d stands for diamagnetic, $i=x,y,z$ are defined with respect to the laboratory frame, sum over repeated indices is implied, $\bar{\boldsymbol{\mu}}$ is the ground state electric dipole moment of the molecule, $\langle\dots\rangle$ stands for thermal average over all orientations, $\langle\dots\rangle_{\rm rot}$ stands for pure rotational average (i.e. isotropic average), σ_{ij} is the nuclear shielding tensor, and σ_{ijk} the shielding polarisability tensor defined as an energy third-derivative at zero-fields:

$$\sigma_{ijk}=\left.\frac{\partial^3 W}{\partial E_i\partial B_j\partial m_k^I}\right|_0. \qquad (2)$$

The pseudoscalar $\sigma^{(1)}$ [5, 9] is evaluated by contraction of $\sigma_{ijk}+\beta\bar{\mu}_i\sigma_{jk}$ with the Levi-Civita totally antisymmetric tensor ϵ_{ijk} [13], leading to $\langle\sigma_{ijk}+\beta\bar{\mu}_i\sigma_{jk}\rangle_{\rm rot}=\sigma^{(1)}\epsilon_{ijk}$, with $\sigma^{(1)}=\frac{1}{6}\epsilon_{abc}\left(\sigma_{abc}+\beta\bar{\mu}_a\sigma_{bc}\right)$, a,b and c defined with respect to the molecule-fixed frame.

In molecules with a degenerate ground state more than one level is populated after the NMR fields partially split the degeneracy. We thus proceed to define a perturbative expansion of the electronic free energy [14] as recently proposed for pNMR chemical shifts [15–17], assuming that electrons are in thermal equilibrium with the NMR fields (i.e. electronic spin-lattice relaxation is fast compared to the NMR dynamics). The electronic free energy $F\left(\boldsymbol{E},\boldsymbol{B},\boldsymbol{m}^I\right)\equiv F$ reads:

$$F=-\beta^{-1}\ln\left\langle \mathrm{Tr}\,\rho\right\rangle_{\mathrm{rot}} \tag{3}$$

where $\rho=e^{-\beta H}$ is the equilibrium statistical operator in the presence of the external fields, fully defined by the molecular Hamiltonian:

$$H=H_0+\lambda V_1+\lambda^2 V_2, \tag{4}$$

$\left\langle \mathrm{Tr}\,\rho\right\rangle_{\mathrm{rot}}$ is the rotationally averaged partition function. Here H_0 is the molecular electrostatic Hamiltonian, V_1 is the sum of the Zeeman V_{zee}, hyperfine V_{hf} and electric dipole/field interaction V_{el} Hamiltonians, and V_2 is the term bilinear in $\boldsymbol{B}$ and $\boldsymbol{m}^I$, responsible for the diamagnetic contribution to the nuclear shielding tensor [18]:

$$V_1=V_{zee}+V_{hf}+V_{el}\quad\text{and}\quad V_2=V_{\mathcal{D}} \tag{5}$$

with

$$\begin{array}{rcl} V_{zee} &=& -\boldsymbol{\mathcal{M}}\cdot\boldsymbol{B}\\ V_{hf} &=& \boldsymbol{\mathcal{F}}\cdot\boldsymbol{m}^I\\ V_{el} &=& -\boldsymbol{\mu}\cdot\boldsymbol{E}\\ V_{\mathcal{D}} &=& \boldsymbol{B}\cdot\boldsymbol{\mathcal{D}}\cdot\boldsymbol{m}^I \end{array} \tag{6}$$

where $\boldsymbol{\mathcal{M}}=\mu_B\left(\boldsymbol{L}+2\boldsymbol{S}\right)$ is the electronic magnetic dipole operator (μ_B the Bohr magneton, in atomic units $\alpha/2$, with α the fine structure constant), $\boldsymbol{\mu}$ is the electric dipole operator, $\boldsymbol{\mathcal{F}}$ is the hyperfine field induced by the electrons at the site of nucleus I, consisting of Fermi contact, spin-dipolar and nuclear spin-electron orbit contributions, and $\boldsymbol{\mathcal{D}}$ is the diamagnetic shielding operator.

The average electric polarization $\overline{P}_i$ is thus defined as:

$$\overline{P}_i=\overline{P}_{d,i}+\overline{P}_{p,i}=-\frac{\partial F}{\partial E_i}, \tag{7}$$

where $\overline{P}_{p,i}$ is the new paramagnetic contribution to $\overline{P}_i$, while $\overline{P}_{d,i}$ follows from Eq. (1). To obtain an analytical expression for Eq. (7) in weak fields, we expand ρ in Eq. (3) in powers of λ:

$$\rho=\rho_0+\lambda\rho_1+\lambda^2\rho_2+\lambda^3\rho_3+\ldots \tag{8}$$

where $\rho_0=e^{-\beta H_0}$, and:

$$\rho_n=-\int_0^\beta d\omega e^{(\omega-\beta)H_0}\left[V_1\rho_{n-1}\left(\omega\right)+V_2\rho_{n-2}\left(\omega\right)\right]. \tag{9}$$

Substituting Eq. (8) into Eq. (3), we obtain:

$$F=F_0+\lambda F_1+\lambda^2 F_2+\lambda^3 F_3+\ldots. \tag{10}$$

Neglecting contributions to $\boldsymbol{P}$ in Eq. (7) that are not bilinear in $\boldsymbol{B}$ and $\boldsymbol{m}^I$, thus could not describe a rotating polarization in the pulsed NMR experiment, and accounting for rotational averaging in the fast spin-lattice relaxation limit, we obtain to leading order in the fields:

$$\overline{P}_i\approx-\frac{\partial F_3}{\partial E_i}=-\Phi_{ijk}B_j m_k^I \tag{11}$$

where we have introduced the *generalized shielding polarizability* Φ_{ijk}, a third-rank tensor defined by:

$$\Phi_{ijk}=\left.\frac{\partial^3 F}{\partial E_i\partial B_j\partial m_k^I}\right|_0=\frac{\partial^3 F_3}{\partial E_i\partial B_j\partial m_k^I}, \tag{12}$$

and where the relevant free-energy contribution F_3 reads:

$$F_3=-\beta^{-1}\left\langle\frac{\mathrm{Tr}\,\rho_3}{\mathrm{Tr}\,\rho_0}\right\rangle_{\mathrm{rot}}. \tag{13}$$

To evaluate Φ_{ijk} we thus first need an explicit expression for F_3 as function of the NMR fields. To obtain $\mathrm{Tr}\,\rho_3$ we use Eq. (9) with $n=3$, leading to:

$$\begin{aligned}\frac{\mathrm{Tr}\,\rho_3}{\mathrm{Tr}\,\rho_0}=&\frac{\beta}{2}\left\langle\int_0^\beta \mathrm{d}u\;T\left[V_1(u)V_2(0)+V_2(u)V_1(0)\right]\right\rangle_0\\ &-\frac{\beta}{6}\left\langle\int_0^\beta\int_0^\beta\int_0^\beta \mathrm{d}u\mathrm{d}v\;T\left[V_1(u)V_1(v)V_1(0)\right]\right\rangle_0,\end{aligned} \tag{14}$$

where $\left\langle O\right\rangle_0\equiv\mathrm{Tr}(\rho_0 O)/\,\mathrm{Tr}\,\rho_0$, the imaginary-time Heisenberg picture operators $V_i(\omega)$ (with $i=1,2$) are:

$$V_i(\omega)=e^{\omega H_0}V_i e^{-\omega H_0}, \tag{15}$$

and $T\left[\ldots\right]$ represents the imaginary-time-ordering operator permuting the $V_i(\omega)$'s so that operators with later imaginary-time arguments are shifted to the left.

We can now evaluate Φ_{ijk} in Eq. (12), by substitution of Eq. (14) into Eq. (13), and by carrying out the third mixed-derivative as prescribed in Eq. (12). We obtain:

$$\begin{aligned}\Phi_{ijk}=&\frac{1}{6}\sum_{P\in S_3}\left\langle\int_0^\beta\int_0^\beta\int_0^\beta \mathrm{d}u\mathrm{d}v\;T\left[P\mu_i(u)\mathcal{M}_j(v)\mathcal{F}_k\right]\right\rangle_{0,\mathrm{rot}}\\ &+\frac{1}{2}\sum_{P\in S_2}\left\langle\int_0^\beta \mathrm{d}u\;T\left[P\mu_i(u)\mathcal{D}_{jk}\right]\right\rangle_{0,\mathrm{rot}}.\end{aligned} \tag{16}$$

The two terms on the right-hand side of Eq. (16) feature a sum over all permutations of the operators involved, with each permutation P switching the operators without moving their imaginary-time arguments.

Eq. (16) provides a generalization of Buckingham's theory to ground states of arbitrary degeneracy, and hence represents a central result of this Letter. Note that

we can identify Φ_{ijk} in Eq. (16) with the shielding polarizability tensor σ_{ijk} in Eq. (1) [3, 5, 10], however here featuring all contributions, both temperature-independent and temperature-dependent, in a single expression. Temperature dependent contributions were previously discussed [9] for diamagnetic molecules, as summarized by Eq. (1). Our result Eq. (16) also includes new contributions that are non-zero only for paramagnetic molecules, essentially arising from the NMR paramagnetic shift $\beta\sigma^p$, whose theory [19–21] has been recently generalized to ground states of arbitrary degeneracy [15–17, 22].

Explicit sum-over-states expressions for Φ_{ijk} can be derived using the basis $H_0 \left|m\mu\right\rangle = \epsilon_m \left|m\mu\right\rangle$, with $\mu = 1, \ldots, d_m$ enumerating degenerate states, by introducing the resolution of identity $I = \sum_m \sum_{\mu=1}^{d_m} \left|m\mu\right\rangle \left\langle m\mu\right|$ in between operators in Eq. (16), thus turning the thermal propagators Eq. (15) into exponential functions of the integration variables.

For the special but common case of a thermally isolated ground state (TIGS), $\left|a\alpha\right\rangle$ ($\alpha = 1, \ldots, d_a$), this leads to:

$$\Phi_{ijk} = \frac{1}{6}\epsilon_{abc}\left(\sigma_{abc} + \beta\bar{\mu}_a\sigma_{bc} + \beta^2\sigma^p_{abc}\right)\epsilon_{ijk} \tag{17}$$

where σ_{ab} and σ_{abc} in Eq. (17) are equivalent to the closed-shell magnetic shielding and shielding polarizability tensors in Eq. (1), which are generally small second and third order response properties [3, 5, 10]. On the other hand, the new paramagnetic contribution in Eqs. (17) features the third rank tensor σ^p_{abc}:

$$\sigma^p_{abc} = d_a^{-1}\sum_{\alpha\alpha'\alpha''}\Re\left[\left\langle a\alpha\right|\mu_a\left|a\alpha'\right\rangle\left\langle a\alpha'\right|\mathcal{M}_b\left|a\alpha''\right\rangle\left\langle a\alpha''\right|\mathcal{F}_c\left|a\alpha\right\rangle\right] \tag{18}$$

Since Eq. (18) is independent of inverse energy gaps, it is expected to give a sizeable contribution to the chiral polarization Eq. (11). This will be assessed next via ab initio calculations.

Estimate of the chiral polarization. It follows immediately from Eqs. (1), (11), (17) and (18) that $\overline{P}_{p,i}$ for an ensemble of tumbling molecules with a TIGS reads:

$$\overline{P}_{p,i} = -\beta^2\sigma^{(1p)}\left(\boldsymbol{B}\times\boldsymbol{m}^I\right)_i \tag{19}$$

where $\sigma^{(1p)} \equiv \frac{1}{6}\epsilon_{abc}\sigma^p_{abc}$. Since $\boldsymbol{m}^I$ is rotating following a $\pi/2$ pulse, the chiral electric polarization $\overline{P}_{p,i}$ follows such rotation, which can be observed as an AC-voltage induced in a capacitor sandwiching the pick-up coil in a modified NMR/pNMR experiment [5, 23].

Our aim in this last part is to provide an estimate of $\beta^2\sigma^{(1p)}$ at room temperature, and determine the voltage that the associated electric polarization Eq. (19) will produce. We consider here the case of a magnetically anisotropic Kramers Doublet (KD) ground state with a non-zero permanent electric dipole moment. As it will become apparent soon, strong magnetic anisotropy is found to be an essential property for a chiral paramagnetic molecule to produce a large rotating electric polarization in a pulsed NMR experiment. This property is commonly encountered in low-symmetry Ln^{3+} complexes displaying single-molecule magnet (SMM) behaviour [24], for which detection of paramagnetic shifts in ^{1}H-NMR and ^{13}C-NMR experiments has proved very useful in the elucidation of their electronic structure [25, 26]. The ground KD can be associated to a pseudo-spin $\tilde{s} = 1/2$, so that the magnetic moment $\mathcal{M}_i$, and hyperfine field $\mathcal{F}_i$ operators, written in terms of the pseudo-spin operator $\tilde{\boldsymbol{S}}$, are $\mathcal{M}_i = -\mu_B g_{ij}\tilde{S}_j$ and $\mathcal{F}_i = \left(g_I\mu_N\right)^{-1}A^I_{ji}\tilde{S}_j$, where g_{ij} and A^I_{ji} are the Zeeman g-tensor and hyperfine A–tensor, respectively, collecting the matrix elements of $\boldsymbol{\mathcal{M}}$ and $\boldsymbol{\mathcal{F}}$ in the basis of e.g. ab initio KD wavefunctions. The representation of the electric moment component μ_i in the KD basis is forced by time-reversal symmetry to be a scalar $\bar{\mu}_i$ times the 2×2 identity matrix. Within this pseudo-spin representation [15], we obtain:

$$\sigma^{(1p)} = -\frac{\mu_B}{24 g_I\mu_N}\epsilon_{abc}\bar{\mu}_a g_{bd}A^I_{dc}. \tag{20}$$

Since $\sigma^{(1p)}$ Eq. (20) is now expressed in terms of the well-known response tensors g_{ik} and A^I_{kj}, we can proceed with its evaluation via ab initio methods. We choose here to avoid accurate evaluation of A^I_{ij}, which is a non trivial task for a KD with strong multiconfigurational character [27, 28], and invoke instead the well-known dipolar approximation [19] expressing A^I_{ij} as the dipolar magnetic field induced at the site of the probed nucleus I by the paramagnetic ion, making it proportional to the g–tensor itself and the inverse cube of distance $\boldsymbol{R}_I$ between the paramagnetic ion and the probed nucleus I:

$$A^I_{ij} \approx \frac{\mu_B\mu_N g_I}{R_I^5}g_{ik}\left(\delta_{kj}R_I^2 - 3R_{I,k}R_{I,j}\right). \tag{21}$$

Introducing polar coordinates for nucleus I with respect to the reference frame in which the g-tensor is diagonal $\mathbf{g} = \mathrm{diag}\left(g_{xx}, g_{yy}, g_{zz}\right)$ (principal axes frame), $X_I = R_I\sin\chi\cos\Omega$, $Y_I = R_I\sin\chi\sin\Omega$, and $Z_I = R_I\cos\chi$, and substituting Eq. (21) into Eq. (20), we obtain:

$$\begin{aligned}\sigma^{(1p)} = \frac{\mu_B^2}{8}\frac{\sin\chi}{R_I^3}\Big[&\overline{\mu}_x\left(g_{yy}^2 - g_{zz}^2\right)\sin\Omega\cos\chi \\ &+\overline{\mu}_y\left(g_{zz}^2 - g_{xx}^2\right)\cos\Omega\cos\chi \\ &+\overline{\mu}_z\left(g_{xx}^2 - g_{yy}^2\right)\sin\chi\frac{\sin 2\Omega}{2}\Big]\end{aligned} \tag{22}$$

Eq. (22) exposes the fundamental role that magnetic anisotropy plays in the physics predicted here. The pseudoscalar $\sigma^{(1p)}$ is in fact proportional to the difference between the squares of the g-tensor principal components. If the system is magnetically isotropic ($g_{zz} \sim g_{xx} \sim g_{yy}$), $\sigma^{(1p)} \sim 0$. If instead it features a large easy-axis magnetic moment M_0 ($g_{zz} \sim 2M_0, g_{xx} \sim g_{yy} \sim 0$), the effect

is maximal, with $\sigma^{(1p)} \sim \frac{\mu_B^2}{4}\frac{\sin 2\chi}{R_I^3} M_0^2 \mu_\perp$, where $\mu_\perp$ is the component of $\bar{\boldsymbol{\mu}}$ perpendicular to the plane defined by the easy-axis and the probed nucleus.

As discussed in [9], in a diamagnetic sample the dominant mechanism responsible for the rotating chiral polarization is related to the temperature-dependent partial orientation of $\bar{\boldsymbol{\mu}}$ under the effect of the rotating field $\boldsymbol{B} \times \boldsymbol{m}^I$. This is driven by the antisymmetric part of the nuclear shielding $\boldsymbol{\sigma}$, which is a generally small second order response property. While also in paramagnetic molecules partial orientation of $\bar{\boldsymbol{\mu}}$ under $\boldsymbol{B} \times \boldsymbol{m}^I$ is the dominant effect, Eq. (22) shows that this is now driven by an altogether different mechanism which is itself temperature dependent (hence the overall β^2 dependence), based on the anisotropy of the first order Zeeman energy of the paramagnet in the strong NMR magnetic field. In other words, the new mechanism found here is based on the tendency of the molecule to partially align its principal magnetic axis (if any) along the external field. This is a *first order effect* and thus a much larger orientational driving force than that active in a diamagnetic sample.

Guided by this insight, in order to estimate $\beta^2\sigma^{(1p)}$ for promising candidate systems, we considered ten of the Dy^{3+} complexes studied in [29], which we label here **1**–**10** (see Supplementary Material (SM) [30] for more information on their chemical structure). In these systems the ground ${}^6H_{\frac{15}{2}}$ spin-orbit multiplet of Dy^{3+} is split by the low-symmetry crystal field into eight KDs, with a strongly anisotropic ground KD consisting of an almost pure $M_J = \pm 15/2$ atomic doublet, thus all excellent candidates to observe the predicted effect. We performed ab initio calculations using Molcas 8.0 [31] to determine the g-tensor principal axes, and the electric dipole moment components with respect to such axes, within the well-known CASSCF/RASSI–SO approximation [32]. Note that in ionic systems (e.g. complex **6**) the electric dipole $\bar{\boldsymbol{\mu}}$ is origin dependent. However, since the observable polarization results from the torque applied to the molecule by the rotating field $\boldsymbol{B} \times \boldsymbol{m}^I$, the detectable $\bar{\boldsymbol{\mu}}$ is that computed with respect to the centre of mass. The results for $\beta^2\sigma^{(1p)}$ at 293K for all nuclei of all complexes using Eq. (22), are reported in Tables XIII–XXII of the SM [30], with details of calculations.

Our largest estimate for protons at 293K is obtained for complex **6** [33], $[Dy(paaH^*)_2(NO_3)_2(MeOH)]^+$ (paaH* = N–(2–pyridyl)acetoacetamide), where for H_{22} we obtain $\beta^2\sigma^{(1p)} = 7.33 \cdot 10^6$ ppm au ($1\,\text{ppm au} = 10^{-6}\, e\, a_0\, E_h^{-1} \sim 2 \cdot 10^{-18}\,\text{m V}^{-1}$). In fact **1**–**10** all display similar easy-axis anisotropy, but **6** also features the largest $\mu_\perp \sim 20\text{D}$. Note that the largest pseudoscalar obtained in closed-shell molecules for protons is $\sigma^{(1)} = 8.75 \cdot 10^2\,\text{ppm au}$ in (2R)-2-methyloxirane [12], which, even assuming a pure chiral liquid, generates an AC-voltage V smaller than the threshold for experimental detection (1μV), and in a dilute solution (0.1M), it generates $V < 1$nV. Our result is 10^4 times larger, and conservatively assuming a 0.1M solution, and an NMR field of 14.1T, we obtain a chiral polarization $P \sim 8.4\,\text{fC m}^{-2}$, and a voltage $V \sim 15.2\mu$V, well above the 1μV threshold (see SM [30] for details of calculation of P and V). Higher solubilities, thus higher P and V, can be achieved with appropriate choice of ligands, e.g. $Na_9[Ln(W_5O_{18})_2]$ SMMs achieve > 0.1M in water [34], and Ln bis-octa(ethyl)tetraazaporphyrin are highly soluble in common organic solvents [35]. In fact, we find that **1**–**10** are all suitable for ^{1}H-NMR chiral discrimination, with maximal P (fC m^{-2}) and V (μV), reported as (P, V): $(1.3, 2.3)$ for H_{24} in **1**, $(1.9, 3.4)$ for H_{11} in **2**, $(1.5, 2.7)$ for H_{15} in **3**, $(1.9, 3.4)$ for H_1 in **4**, $(2.8, 5.0)$ for H_8 in **5**, $(1.8, 3.3)$ for H_{22} in **7**, $(3.4, 6.1)$ for H_{16} in **8**, $(5.2, 9.3)$ for H_{20} in **9**, and $(4.2, 7.6)$ for H_{18} in **10**. Considering more exotic nuclei, the highest value for a diamagnetic system was obtained for ^{125}Te [23], with $\sigma^{(1)} = 8.28 \cdot 10^4\,\text{ppm au}$ (in a 0.1M solution, $V \sim 18$ nV), while our best result is $\beta^2\sigma^{(1p)} = 2.74 \cdot 10^7\,\text{ppm au}$ for ^{17}O (O_1 in complex **6**).

We conclude by noting that while 3d-complexes are generally less anisotropic than 4f systems, low-coordination environments can trigger unquenched orbital angular momentum, as recently discussed for SMMs based on e.g. Fe^+ [36]. Suitable candidates for detection of the chiral polarization could also be e.g. chiral versions of trigonal bis-trispyrazolylborate chelates of Co^{2+} having $g_\parallel^2 - g_\perp^2 \sim 71$ [37–39], only 5.6 times smaller than in the Dy^{3+} complexes shown here, where $g_\parallel^2 - g_\perp^2 \sim 400$. Since our proposal specifically addresses chiral molecules with strong magnetic anisotropy, it could provide an alternative route to *direct* chiral discrimination *in solution* for these specific systems when other techniques such as circular dichroism are not viable (e.g. if ligands have no chromophores). Despite the specificity of the proposed approach, we note that synthesis and characterization of enantiomerically pure magnetically anisotropic complexes is strategic in several research areas, among which enantioselective catalysis [40], indirect chiral discrimination via NMR using chiral Ln^{3+} shift reagents [40], and stereoselective binding of DNA chains aimed at inhibiting particular biological functions [41]. We further note that the method could extend the current scope of Ln^{3+} shift reagents, to include *direct* chiral discrimination of chiral analytes, via the use of *achiral* Ln^{3+} shift reagents.

We hope these results will encourage experimentalists to gauge our predictions, and design the first experiment to achieve chiral discrimination in pNMR spectroscopy.

We thank Dr C. Boskovic and Dr M. Vonci (University of Melbourne), Prof. K. S. Murray and Prof. G. B. Deacon (Monash University), for useful discussions on the solubility of Ln-SMMs. A.S. acknowledges support from the Australian Research Council, grant ID: DP150103254

* asoncini@unimelb.edu.au

[1] K. W. Busch and M. A. Busch, *Chiral Analysis* (Elsevier, Amsterdam, 2006).
[2] P. Fischer, D. S. Wiersma, R. Righini, B. Champagne, and A. D. Buckingham, Phys. Rev. Lett. **85**, 4253 (2000).
[3] A. D. Buckingham, Chem. Phys. Lett. **398**, 1 (2004).
[4] P. Fischer and F. Hache, Chirality **17**, 421 (2005).
[5] A. D. Buckingham and P. Fischer, Chem. Phys. **324**, 111 (2006).
[6] Y. Li and C. Bruder, Phys. Rev. A **77**, 015403 (2008).
[7] K. Hirmatsu, M. Okuno, H. Kano, P. Leproux, V. Couderc, and H.-o. Hamaguchi, Phys. Rev. Lett. **109**, 083901 (2012).
[8] D. Patterson, M. Schnell, and J. M. Doyle, Nature **497**, 475 (2013).
[9] A. D. Buckingham, J. Chem. Phys. **140**, 011103 (2014).
[10] P. Lazzeretti, A. Soncini, and R. Zanasi, Theor. Chem. Account **119**, 99 (2008).
[11] G. Monaco and R. Zanasi, Chirality **23**, 752 (2011).
[12] A. D. Buckingham, P. Lazzeretti, and S. Pelloni, Mol. Phys. **113**, 1780 (2015).
[13] D. P. Craig and Thirunamachandran, *Molecular Quantum Electrodynamics* (Academic Press, London, 1984) see Appendix 2.
[14] R. P. Feynman, *Statistical Mechanics: A Set of Lectures* (Benjamin, Reading, MA, 1972) see sections 2.10 and 2.11, pp. 66–71.
[15] W. Van den Heuvel and A. Soncini, Phys. Rev. Lett. **109**, 073001 (2012).
[16] A. Soncini and W. Van den Heuvel, J. Chem. Phys. **138**, 021103 (2013).
[17] W. Van den Heuvel and A. Soncini, J. Chem. Phys. **138**, 054113 (2013).
[18] A. Abragam, *The principles of Nuclear Magnetism* (Oxford University Press, Oxford, 1961).
[19] R. J. Kurland and B. R. McGarvey, J. Magn. Reson. **2**, 286 (1970).
[20] S. Moon and S. Patchkovskii, in *Calculation of NMR and EPR Parameters. Theory and Applications*, edited by M. B. M. Kaupp and V. G. Malkin (Wiley-VCH Verlag GmbH, Weinheim, 2004) Chap. 20.
[21] T. O. Pennanen and J. Vaara, Phys. Rev. Lett. **100**, 133002 (2008).
[22] B. Martin and J. Autschbach, J. Chem. Phys. **142**, 054108 (2015).
[23] S. Pelloni, F. Faglioni, and P. Lazzeretti, Rend. Fis. Acc. Lincei **24**, 283 (2013).
[24] D. N. Woodruff, R. E. P. Winpenny, and R. A. Layfield, Chem. Rev. **113**, 5110 (2013).
[25] N. Ishikawa, M. Sugita, T. Okubo, N. Tanaka, T. Iino, and Y. Kaizu, Inorg. Chem. **42**, 2440 (2003).
[26] M. Damjanovic, T. Morita, K. Katoh, M. Yamashita, and M. Enders, Chem. Eur. J. **21**, 14421 (2015).
[27] F. Aquino, B. Pritchard, and J. Autschbach, J. Chem. Theory and Comput. **8**, 598 (2012).
[28] K. Sharkas, B. Pritchard, and J. Autschbach, J. Chem. Theory and Comput. **11**, 538 (2015).
[29] N. F. Chilton, D. Collison, E. J. L. McInnes, R. E. P. Winpenny, and A. Soncini, Nat. Commun. **4**, 2551 (2013).
[30] For more information on EPAPS, see http://www.aip.org/pubservs/epaps.html.
[31] F. Aquilante, L. De Vico, N. Ferré, G. Ghigo, P.-Å. Malmqvist, P. Neogrády, T. B. Pedersen, M. Pitonak, M. Reiher, B. O. Roos, L. Serrano-Andrés, M. Urban, V. Veryazov, and R. Lindh, J. Comput. Chem. **31**, 224 (2010).
[32] L. Ungur and L. F. Chibotaru, in *Lanthanides and Actinides in Molecular Magnetism*, edited by R. A. Layfield and M. Murugesu (Wiley–VCH, 2015) pp. 153–184.
[33] N. F. Chilton, S. K. Langley, B. Moubaraki, A. Soncini, S. R. Batten, and K. S. Murray, Chem. Sci. **4**, 1719 (2013).
[34] C. Boskovic and M. Vonci, (private communication).
[35] N. Gimenez-Agullo, C. Saenz de Pipaon, L. Adriaenssens, M. Filibian, M. Martinez-Belmonte, E. C. Escudero-Adan, P. Carretta, P. Ballester, and J. R. Galan-Mascaros, Chem. Eur. J. **20**, 12817 (2014).
[36] J. M. Zadrozny, D. J. Xiao, M. Atanasov, G. J. Long, F. Grandjean, F. Neese, and J. R. Long, Nature Chemistry **5**, 577 (2013).
[37] V. V. Novikov, A. A. Pavlov, A. S. Belov, A. V. Vologzhanina, A. Savitsky, and Y. Z. Voloshin, J. Phys. Chem. Lett. **5**, 3799 (2014).
[38] D. L. Tierney, J. Phys. Chem. A **116**, 10959 (2012).
[39] J. P. Jesson, J. Chem. Phys. **45**, 1049 (1966).
[40] H. C. Aspinall, Chem. Rev. **102**, 1807 (2002).
[41] C. Zhao, J. Ren, J. Gregoliński, J. Lisowski, and X. Qu, Nucl. Acids Res. **40**, 8186 (2012).

arXiv:1604.01626v1 [physics.chem-ph] 6 Apr 2016

Room temperature chiral discrimination in paramagnetic NMR spectroscopy: Supplementary Material

Alessandro Soncini* and Simone Calvello

School of Chemistry, University of Melbourne, VIC3010, Australia

Computational details

All calculations were performed with the MOLCAS-8.0 program [1].

Computational model. As probe molecules, we chose ten Dy^{3+} complexes with strongly anisotropic easy-axis ground states (see Table I for names, using the experimental geometries from Ref. [2] and reported below in Tables III–XII. Crystallographic data for these molecules show that they do not possess symmetry (C_1 group).

	Compound
1	$[Dy(acac)_3(H_2O)_2]$
2	$[Dy(acac)_3(dppz)]$
3	$[Dy(acac)_3(dpq)]$
4	$[Dy(acac)_3(phen)]$
5	$[Dy(hfac)_3(dme)]$
6	$[Dy(paaH^*)_2(NO_3)_2(MeOH)]^+$
7	$[Dy(tfpb)_3(dppz)]$
8	$[Dy(tta)_3(bipy)]$
9	$[Dy(tta)_3(phen)]$
10	$[Dy(tta)_3(pinene\text{-}bipy)]$

TABLE I: The ten Dy complexes **1**–**10** for which ab initio calculation of the chiral polarization was carried out. Acronyms used above: acac = acetylacetonate, dppz = dipyridophenazine, dpq = dipyridoquinoxaline, phen = 1,10-phenanthroline, hfac = hexafluoroacetylacetone, dme = dimethoxyethane, paaH* = N-(2-Pyridyl)acetoacetamide, tfpb = 4,4,4-trifluoro-1-phenyl-1,3-butandionate, tta = tetradecylthioacetate, bipy = 2,20-bipyridine, pinene-bipy = 4,5-pinene bipyridine.

Basis sets. ANO-rcc basis sets from the MOLCAS-8 basis set library are used with the following contractions: VTZP on Dy, VTZ on N and O and VDZ on other nuclei.

CASSCF. The active space consists of the seven valence $4f$ orbitals of the Dy^{3+} ion, occupied by 9 electrons. For a given spin, CASSCF calculations are averaged over all the roots that derive from

the $4f^n$ electron configuration. For Dy^{3+} we optimised the average energy of the 21 CASSCF roots with $S = 5/2$ (from the free ion terms ^{6}H, ^{6}F, ^{6}P), in accordance with previous calculations [2].

RASSI. In this step the spin-orbit coupling Hamiltonian was diagonalized in the basis of the optimized CASSCF wave functions to yield the twofold degenerate ground state. Electric dipole components were also calculated. The module SINGLE_ANISO [3] has been used to compute the principal magnetic axes and related g-tensor principal components.

All calculations have been performed using the Dysprosium atom as the origin of the coordinate system. Since the rotating electric polarization formula requires all properties to be calculated at the center of mass, the electric dipole moment $\boldsymbol{\mu}$ (expressed in a.u.) calculated in the former coordinates has to be translated into the center of mass using the formula:

$$\boldsymbol{\mu'} = \boldsymbol{\mu} - Q\mathbf{C}_{\mathbf{mass}} \tag{1}$$

where Q is the total charge of the molecule and $\mathbf{C}_{\mathbf{mass}}$ is the coordinate of the center of mass expressed in atomic units.

Compound	CM Coordinates			Dipole Moment		
	x	y	z	x	y	z
1	-0.735	0.588	-0.078	1.997	-0.756	0.226
2	2.058	-1.826	-1.538	1.173	-1.536	-1.426
3	1.425	1.790	0.547	1.726	2.182	0.433
4	0.937	-0.264	0.860	2.188	-0.409	2.132
5	-0.687	-0.379	0.690	3.442	2.597	-2.092
6	0.611	-0.442	0.147	7.511	-0.739	-1.755
7	0.072	-0.913	0.174	-1.280	-2.357	-1.497
8	0.632	-0.032	0.371	5.925	-1.077	-3.770
9	-0.662	-0.239	0.219	-7.391	0.034	-0.735
10	-0.871	0.414	-0.310	-1.532	7.161	-1.903

TABLE II: Center of mass (CM) coordinates, in a.u., and dipole moment $\boldsymbol{\mu'}$, expressed in a.u., translated (for the ionic molecule **6**) in the center of mass for the studied compounds.

Pseudoscalar and voltage

Using Eq. P22, the paramagnetic contribution to the pseudoscalar, $\beta^2\sigma^{(1p)}$, calculated at T = 293 K, was computed for all non-lanthanide atoms of the ten probe complexes, and reported below

in Tables XIII–XXII. The rotating polarization and the induced voltage were calculated using the following formulae [4, 5]:

$$V^I = P_y^I \frac{d}{(\epsilon - 1)\epsilon_0} = \frac{N^I(\beta^2\sigma^{(1p)})(\hbar\gamma_I B_z)^2 I^I(I^I+1)}{3kT}\frac{d}{(\epsilon-1)\epsilon_0} \tag{2}$$

where ϵ is the dielectric constant of the medium, ϵ_0 is the permittivity of free space, and d is the separation between the two plates of the capacitor sandwiching the NMR pick-up coil for the chiral polarization detection. For the nucleus I γ_I is the gyromagnetic ratio, I^I is the nuclear spin and $N^I = M \cdot NA$, where M is the molarity and NA is Avogadro's number, is the number density, approximable to $N^I \approx 10^{28}\mathrm{m}^{-3}$ for a pure chiral liquid [4]. We assumed a 0.1M solution of our complex in an appropriate solvent, as a conservative estimate of a suitable concentration obtainable in practice (see text for further comments on solubility), yielding a number density of $N^I = 6.022 \cdot 10^{25}\mathrm{m}^{-3}$. In the calculations performed, the ratio $\frac{d}{(\epsilon-1)}$ was set equal to 0.016 m and B_z was set to 14.1 T, as suggested in previous works [4, 5].

The highest value of $\beta^2\sigma^{(1p)}$ obtained for the hydrogen atoms, which constitute the easiest and the most frequently measured NMR nucleus, consists of $7.33 \cdot 10^6$ ppm au for H_{22} in complex **6**, 10^4 times larger than the largest value obtained for diamagnetic systems, in (2R)-2-dimethyloxirane [6]. Including all the nuclei, the highest value increases to $2.74 \cdot 10^7$ ppm au for an oxygen atom (O_1) in complex **6**, to be compared with the highest previous estimate of $8.28 \cdot 10^4$ ppm au for a tellurium nucleus in 1,2-M-ditellurin [6].

The maximum voltage we obtained consists of 15.2 μV, induced by the resonant precession of H_{22} in complex **6** (see table XVIII), well above the experimental threshold of 1μV.

GEOMETRIES

Atom	x	y	z	Atom	x	y	z	Atom	x	y	z
Dy	0.00000	0.00000	0.00000	C1	-1.46448	-3.94277	2.05137	H1	-0.68899	-4.00038	2.61444
O1	-0.66534	-1.90736	1.18518	C2	-1.17322	-3.02569	0.88540	H2	-1.68698	-4.81693	1.72281
O2	-1.06398	-1.42002	-1.51292	C3	-1.50151	-3.43635	-0.39120	H3	-2.20169	-3.59310	2.55691
O3	-0.76256	0.82638	2.09716	C4	-1.44490	-2.62318	-1.51845	H4	-1.78026	-4.31627	-0.50514
O4	-2.02054	1.11207	-0.34927	C5	-1.87094	-3.19417	-2.85478	H5	-2.74477	-2.86794	-3.07935
O5	1.08251	2.00888	0.37000	C6	-1.15415	2.47835	3.73505	H6	-1.89110	-4.15336	-2.80168
O6	0.21421	1.09558	-2.10487	C7	-1.37365	1.89584	2.37107	H7	-1.24523	-2.92556	-3.53180
O7	1.79374	-0.47691	1.57270	C8	-2.23926	2.54748	1.49714	H8	-1.52805	1.89226	4.39769
O8	1.77346	-1.16850	-1.08149	C9	-2.56018	2.09511	0.22276	H9	-1.57997	3.33661	3.78926
				C10	-3.64352	2.82205	-0.55492	H10	-0.21169	2.57768	3.89214
				C11	1.45860	4.32104	0.63649	H11	-2.62944	3.34059	1.78476
				C12	1.03189	3.12781	-0.18986	H12	-3.37471	2.90947	-1.47199
				C13	0.62121	3.33761	-1.51071	H13	-3.77964	3.69224	-0.17548
				C14	0.25490	2.33232	-2.38463	H14	-4.46189	2.32139	-0.50846
				C15	-0.07738	2.70007	-3.80504	H15	0.98537	4.31806	1.47170
								H16	1.26005	5.12864	0.15749
								H17	2.40280	4.27236	0.80353
								H18	0.59174	4.21276	-1.82045
								H19	0.62128	2.39093	-4.38692
								H20	-0.15758	3.65449	-3.87805
								H21	-0.90759	2.28961	-4.05615
								H22	2.09150	-1.30637	1.90977
								H23	1.69868	-1.55472	-1.95099
								H24	1.88696	0.13997	2.24164
								H25	2.66661	-1.27856	-0.83369

TABLE III: Experimental geometry, used in calculations, for complex **1**. All coordinates are expressed in Å.

Atom	x	y	z	Atom	x	y	z	Atom	x	y	z
Dy	0.00000	0.00000	0.00000	C1	1.93717	-2.65581	-1.17994	H1	5.88651	-7.12522	-2.60195
O1	-1.32985	1.41244	1.21864	C2	5.96298	-6.50784	-3.29283	H2	7.49440	-5.97818	-6.08490
O2	0.17925	-0.63848	2.21859	C3	5.19834	-5.35245	-3.27205	H3	-2.80444	2.10705	-2.99414
O3	2.19456	0.39502	0.61607	C4	2.60130	-3.78727	-1.21760	H4	-4.14306	1.29725	-2.90324
O4	-1.36991	0.44785	-1.80848	C5	6.93650	-5.80512	-5.36026	H5	-3.03104	0.86254	-3.91877
O5	0.51316	2.17971	-0.71113	C6	-3.20044	1.23820	-3.04998	H6	6.37014	-4.09271	-6.00698
O6	-1.83849	-1.41503	0.16804	C7	2.83080	0.39631	1.68226	H7	3.66050	-0.97368	-4.94210
N1	1.04131	-2.33454	-0.19038	C8	0.97764	-0.40450	3.09258	H8	2.77882	-5.61325	-0.27869
N2	1.38791	-0.49107	-2.08639	C9	-0.14429	3.24699	-0.74360	H9	2.82112	0.17596	3.70294
N3	4.32928	-5.18625	-2.26950	C10	6.25165	-4.70347	-5.31481	H10	2.46858	0.94822	-4.72782
N4	4.65260	-3.26447	-4.25382	C11	3.82084	-3.08899	-3.26036	H11	1.02676	1.18226	-2.97986
				C12	3.02033	-1.83005	-3.22920	H12	7.34870	-7.52497	-4.34473
				C13	3.10494	-0.84880	-4.20707	H13	-1.67225	4.24868	-0.09948
				C14	2.34351	-4.79263	-0.25661	H14	1.22541	-5.12877	1.34331
				C15	-3.00231	-1.35556	-0.23843	H15	-2.84152	2.48644	2.69909
				C16	3.61088	-4.05303	-2.26820	H16	-3.49293	3.37562	1.58486
				C17	2.13509	-1.59265	-2.20847	H17	-2.30259	3.93579	2.43677
				C18	2.29085	0.07129	2.94583	H18	4.59173	0.81811	0.79529
				C19	5.35924	-4.37384	-4.28888	H19	4.74608	0.43949	2.30950
				C20	2.38929	0.27061	-4.09539	H20	4.20478	1.85404	1.90692
				C21	-2.58465	0.33394	-1.99029	H21	-0.25470	-0.12628	4.70289
				C22	-1.68711	2.60056	1.01346	H22	1.22404	-0.43124	5.12495
				C23	1.53521	0.40630	-3.03310	H23	0.29935	-1.59026	4.61588
				C24	6.83296	-6.74882	-4.32914	H24	0.21776	-3.12530	1.39006
				C25	-1.20389	3.45543	0.03039	H25	-3.61851	-2.84975	0.99398
				C26	1.43438	-4.49707	0.69269	H26	-4.48001	-2.74671	-0.31115
				C27	-2.67105	3.14973	2.02640	H27	-4.68281	-1.70791	0.84593
				C28	4.21487	0.92401	1.67317	H28	-4.30879	-0.44326	-1.42798
				C29	0.51538	-0.66146	4.51459	H29	1.27219	4.33623	-1.71368
				C30	0.82728	-3.29608	0.70828	H30	-0.04749	5.14400	-1.46175
				C31	-4.04167	-2.24539	0.37842	H31	0.00458	4.06350	-2.59676
				C32	-3.39771	-0.48301	-1.24358				
				C33	0.30930	4.29138	-1.71628				

TABLE IV: Experimental geometry, used in calculations, for complex **2**. All coordinates are expressed in Å.

Atom	x	y	z	Atom	x	y	z	Atom	x	y	z
Dy	0.00000	0.00000	0.00000	C1	3.40693	0.06819	0.68587	H1	3.26932	-0.83997	0.53907
O1	0.77990	1.00205	-1.93607	C2	4.68842	0.50894	0.95107	H2	5.39802	-0.09191	0.94475
O2	1.36124	-1.55616	-1.03631	C3	4.90786	1.80228	1.21468	H3	5.77029	2.10488	1.38990
O3	-1.38343	-1.79498	-0.39226	C4	3.82396	2.70016	1.22731	H4	6.08778	6.16473	2.54697
O4	-1.91156	0.85823	-1.01421	C5	3.97367	4.09205	1.57143	H5	4.28069	7.53053	2.46331
O5	-1.40924	0.43114	1.85715	C6	5.27268	5.80920	2.27388	H6	0.52778	6.18288	1.14207
O6	0.82386	-1.21070	1.78927	C7	4.16817	6.64152	2.21548	H7	-1.49287	5.26496	0.46645
N1	2.35836	0.89240	0.63062	C8	2.88986	4.95384	1.51145	H8	-1.63913	3.00845	0.21704
N2	0.21794	2.52477	0.51539	C9	1.58360	4.43840	1.09945	H9	-2.71876	1.63539	3.55250
N3	5.21296	4.53326	1.95817	C10	0.45758	5.27137	0.97316	H10	-2.07978	0.97414	4.82164
N4	2.98428	6.26244	1.83347	C11	-0.74391	4.72929	0.59747	H11	-3.16336	0.19849	3.99607
				C12	-0.80783	3.37151	0.42068	H12	-0.34818	-0.62451	4.64011
				C13	1.41647	3.09084	0.83899	H13	1.73351	-2.80596	3.66773
				C14	2.56462	2.19886	0.88950	H14	1.61982	-1.70510	4.77745
				C15	-2.42401	0.80821	3.94082	H15	2.62783	-1.52068	3.59196
				C16	-1.35440	0.21455	3.09314	H16	3.61406	-2.87402	-1.69140
				C17	-0.32888	-0.54311	3.71509	H17	3.14215	-2.86796	-3.18628
				C18	0.68879	-1.16128	3.03632	H18	2.28373	-3.59855	-2.09709
				C19	1.76676	-1.86273	3.84295	H19	2.87197	-0.63547	-3.62353
				C20	2.84976	-2.84284	-2.27230	H20	-3.69058	-1.27273	-2.47594
				C21	2.09790	-1.59705	-2.03079	H21	1.16565	2.03112	-4.29126
				C22	2.27923	-0.51939	-2.91635	H22	2.61523	1.45499	-4.44280
				C23	1.63758	0.69852	-2.81059	H23	2.31564	2.54994	-3.36150
				C24	-2.35912	-1.99624	-1.19100	H24	-3.87555	2.10174	-1.76244
				C25	-3.04258	-1.00698	-1.86346	H25	-4.62622	0.92662	-2.48067
				C26	-2.83943	0.35384	-1.69772	H26	-3.40832	1.62277	-3.17996
				C27	1.96387	1.78355	-3.81927	H27	-2.81447	-3.61650	-2.34965
				C28	-3.77135	1.33993	-2.33860	H28	-3.55645	-3.63674	-0.96843
				C29	-2.72480	-3.44689	-1.40884	H29	-2.03498	-4.00840	-1.04736

TABLE V: Experimental geometry, used in calculations, for complex **3**. All coordinates are expressed in Å.

Atom	x	y	z	Atom	x	y	z	Atom	x	y	z
Dy	0.00000	0.00000	0.00000	C1	3.36302	0.71987	-0.00923	H1	3.07425	1.24945	-0.71571
O1	0.56613	2.24850	0.10270	C2	4.73756	0.63161	0.23499	H2	5.34397	1.08764	-0.30432
O2	-1.28528	0.93065	1.67658	C3	5.16680	-0.12283	1.25400	H3	6.07936	-0.20269	1.41681
O3	0.98498	0.39393	-2.06179	C4	4.25485	-0.78270	2.07241	H4	5.52754	-1.67163	3.41559
O4	-1.66828	0.87854	-1.37712	C5	4.62529	-1.56446	3.21498	H5	3.99020	-2.62992	4.74278
O5	0.99363	-2.02406	-0.59215	C6	3.71364	-2.14447	3.99850	H6	1.53730	-3.05652	5.27918
O6	-1.65046	-1.63234	0.10270	C7	2.31028	-2.03309	3.70923	H7	-0.65229	-2.72028	4.77766
N1	2.45574	0.09089	0.71033	C8	1.30454	-2.56897	4.52327	H8	-1.21437	-1.56866	2.85012
N2	0.59543	-1.14122	2.24685	C9	0.01458	-2.38404	4.22382	H9	1.30769	4.88715	0.85570
				C10	-0.31508	-1.67374	3.06235	H10	-0.04905	5.24020	0.15503
				C11	1.89174	-1.29967	2.58119	H11	1.07485	4.49417	-0.64448
				C12	2.88948	-0.65472	1.75842	H12	-1.16638	4.05495	1.98664
				C13	0.65659	4.59294	0.21463	H13	-3.20572	1.64874	3.13939
				C14	0.08553	3.26689	0.65073	H14	-2.67379	3.08406	3.47664
				C15	-0.89034	3.23957	1.63340	H15	-1.97643	1.82316	4.09735
				C16	-1.48114	2.10266	2.12619	H16	2.36305	2.38216	-3.50964
				C17	-2.42063	2.16991	3.31965	H17	1.59322	1.80425	-4.74670
				C18	1.91488	1.61091	-3.86433	H18	2.52778	0.87328	-3.90067
				C19	0.74070	1.24945	-2.96597	H19	-0.56158	2.45781	-3.86142
				C20	-0.48536	1.85468	-3.15930	H20	-3.56898	1.69917	-3.07063
				C21	-1.60760	1.62772	-2.38596	H21	-2.77084	3.00000	-3.42678
				C22	-2.93551	2.34223	-2.74647	H22	-3.29062	2.77304	-1.96440
				C23	1.85695	-4.21025	-0.50202	H23	2.08435	-4.33003	-1.42800
				C24	0.71893	-3.24356	-0.39009	H24	1.59994	-5.05505	-0.12407
				C25	-0.56928	-3.69328	-0.08191	H25	2.61712	-3.86560	-0.02813
				C26	-1.66773	-2.90311	0.09543	H26	-0.68718	-4.61163	0.00967
				C27	-3.01622	-3.54827	0.29022	H27	-3.28085	-3.46421	1.20894
								H28	-2.96308	-4.47714	0.05619
								H29	-3.66328	-3.11326	-0.26943

TABLE VI: Experimental geometry, used in calculations, for complex **4**. All coordinates are expressed in Å.

Atom	x	y	z	Atom	x	y	z	Atom	x	y	z
Dy	0.00000	0.00000	0.00000	C1	3.34275	-0.50471	0.57232	H1	3.46536	-1.34257	0.08135
O1	2.39513	0.32214	-0.12253	C2	2.88909	0.78989	-1.38533	H2	3.00644	-0.70000	1.47106
O2	0.71885	1.75691	-1.55593	C3	2.14816	1.99472	-1.72028	H3	4.19928	-0.03537	0.63890
O3	-1.02399	1.94275	0.75540	C4	-0.02383	2.92236	-1.87215	H4	2.75421	0.10007	-2.08227
O4	0.73812	0.39143	2.18463	C5	-1.20616	2.46721	1.87465	H5	3.85772	0.98834	-1.32084
O5	-1.72638	-0.85907	1.31503	C6	-0.64389	2.10182	3.08337	H6	2.33908	2.25459	-2.65646
O6	0.69783	-2.08280	0.76788	C7	0.30197	1.07811	3.13954	H7	2.43367	2.73652	-1.12944
O7	-1.77017	0.15204	-1.49976	C8	-2.15552	3.65471	1.84345	H8	0.15659	3.61534	-1.20434
O8	0.57697	-1.24965	-1.88255	C9	0.89752	0.72690	4.48555	H9	-0.98020	2.70817	-1.87215
F1	-3.33960	3.32712	1.40656	C10	-1.73339	-1.69379	2.25953	H10	0.23716	3.24680	-2.75840
F2	-2.32017	4.23586	3.00639	C11	-0.75074	-2.62458	2.57575	H11	-0.90138	2.54910	3.88016
F3	-1.75616	4.58865	1.03417	C12	0.38079	-2.76002	1.77479	H12	-0.85233	-3.17265	3.34342
F4	2.24625	0.73320	4.45643	C13	-2.95776	-1.61031	3.14786	H13	-1.27622	-0.97719	-4.44768
F5	0.60675	-0.49998	4.87043	C14	1.38796	-3.84988	2.13262				
F6	0.56822	1.54902	5.47999	C15	-1.91205	-0.09050	-2.72511				
F7	-3.08562	-0.43856	3.69708	C16	-1.05026	-0.83229	-3.53647				
F8	-2.97527	-2.55370	4.10692	C17	0.10754	-1.35045	-3.04341				
F9	-4.06126	-1.84026	2.45092	C18	-3.17846	0.47491	-3.35755				
F10	2.56854	-3.45614	2.00780	C19	0.99561	-2.15524	-3.99208				
F11	1.24784	-4.84997	1.37328								
F12	1.26886	-4.34441	3.32261								
F13	-3.69518	1.44822	-2.70847								
F14	-4.13483	-0.55510	-3.07670								
F15	-3.16970	0.62138	-4.64324								
F16	0.99385	-3.43095	-3.60928								
F17	2.21122	-1.74891	-4.00040								
F18	0.61551	-2.16469	-5.23615								

TABLE VII: Experimental geometry, used in calculations, for complex **5**. All coordinates are expressed in Å.

Atom	x	y	z	Atom	x	y	z	Atom	x	y	z
Dy	0.00000	0.00000	0.00000	C1	-0.92578	3.18610	-0.43022	H1	0.88620	-5.13818	-0.48570
O1	0.22937	-2.16529	-0.76376	C2	-2.05447	4.07625	-0.91653	H2	-1.84580	4.40676	-1.81520
O2	1.30069	1.78059	0.79342	C3	1.22593	3.03949	0.72825	H3	-2.15844	4.83521	-0.30533
O3	-1.06221	1.94542	-0.60636	C4	0.89333	-3.18263	-0.38689	H4	-2.88806	3.56165	-0.94311
O4	1.42870	0.40897	-1.84273	C5	4.25215	4.16407	2.58473	H5	4.14321	5.10500	2.51207
O5	-2.44002	-0.47744	-0.15612	C6	3.90801	-2.03931	1.62108	H6	4.54196	-1.42120	1.20104
O6	2.04553	-0.80958	0.83064	C7	2.59018	-1.96520	0.90797	H7	4.25794	-2.95309	1.56669
O7	-1.41215	-0.44868	-2.03796	C8	3.34022	3.29633	1.97089	H8	3.78498	-1.79206	2.56137
O8	-0.96428	0.51283	2.17926	C9	2.06385	-3.13649	0.38237	H9	2.52168	-3.95118	0.55319
O9	-0.68997	-1.58728	1.80266	C10	5.31338	3.63592	3.29809	H10	5.93045	4.21621	3.72819
O10	-1.84589	-0.95427	3.51911	C11	0.17902	3.79082	0.16590	H11	0.22627	4.73926	0.19316
O11	-3.59070	-0.50927	-2.00684	C12	-0.64311	-4.69261	-1.58396	H12	-0.64505	-6.73431	-1.42360
N1	0.43754	-4.44101	-0.78079	C13	-1.07509	-5.99395	-1.83523	H13	6.20871	1.88406	3.87454
N2	2.26752	3.77034	1.28499	C14	5.48030	2.25158	3.38787	H14	4.69778	0.50395	2.83366
N3	3.54373	1.95876	2.06656	C15	4.59217	1.44614	2.77354	H15	0.92435	2.09806	-2.55629
N4	-1.19182	-0.68945	2.53221	C16	1.45587	1.31827	-2.82051	H16	1.08139	0.92964	-3.63852
N5	-2.51153	-0.46552	-1.41876	C17	-2.32704	-3.84945	-3.02971	H17	2.38182	1.59434	-2.98421
N6	-1.29389	-3.66331	-2.17817	C18	-2.76737	-5.10651	-3.30329	H18	-2.74614	-3.10084	-3.43769
				C19	-2.13176	-6.18582	-2.68711	H19	-3.49218	-5.24724	-3.90109
								H20	-2.43675	-7.06853	-2.86115
								H21	2.24482	4.64613	1.29277
								H22	-0.94041	-2.86333	-2.07528
								H23	2.94454	1.43965	1.68282
								H24	2.18804	0.16215	-1.58740

TABLE VIII: Experimental geometry, used in calculations, for complex **6**. All coordinates are expressed in Å.

Atom	x	y	z	Atom	x	y	z	Atom	x	y	z
Dy	0.00000	0.00000	0.00000	C1	0.95173	-3.14104	-0.96504	H1	-2.06903	3.76571	0.88426
O1	-0.24787	2.22212	0.52584	C2	1.72933	-4.21655	-1.40402	H2	-2.65413	-5.63267	0.07636
O2	-1.49778	0.08861	1.78206	C3	-0.39485	-3.36208	-0.46536	H3	-1.93618	2.25066	-3.58489
O3	0.42373	0.71555	-2.17923	C4	-0.91946	-4.65184	-0.38713	H4	-3.86108	-3.76995	0.61691
O4	2.12505	0.91313	0.14174	C5	-0.19218	1.37075	-3.05602	H5	3.18558	-8.80409	-2.17806
O5	-2.05235	0.43826	-1.00590	C6	2.55243	1.96518	0.64843	H6	-1.17301	-9.52769	-0.55409
O6	0.98339	-1.04767	1.82643	C7	-0.08195	-5.79293	-0.76657	H7	-3.29294	5.47297	1.81475
N1	-1.11987	-2.26932	-0.13729	C8	-1.64222	4.39604	1.42014	H8	-0.36743	-0.57592	4.63424
N2	1.36781	-1.86597	-1.04793	C9	-2.26407	-4.79035	0.02499	H9	-0.01976	6.77885	3.61152
N3	-0.60910	-7.02561	-0.62530	C10	-2.65131	0.51232	3.76329	H10	-2.40257	-0.45720	5.52620
N4	2.00520	-6.58381	-1.64219	C11	1.22563	-5.57048	-1.27326	H11	2.29803	3.74593	1.52288
F1	-4.16042	2.48244	-2.85171	C12	-0.27698	4.29146	1.63029	H12	2.86052	-0.76247	-1.66787
F2	-4.60560	0.61577	-2.05548	C13	-1.56025	1.68818	-2.94744	H13	-0.93275	2.18000	-5.49490
F3	-4.13187	2.21392	-0.82144	C14	-2.97636	-3.68798	0.34255	H14	3.49085	-4.65184	-2.32050
F4	1.89892	-1.20172	5.15728	C15	0.59078	1.80887	-4.24102	H15	-2.90005	-1.70372	0.48732
F5	4.55591	3.20322	0.94847	C16	2.31782	-8.93694	-1.86284	H16	-2.23863	7.01345	3.16554
F6	4.64975	1.58642	-0.30308	C17	0.44663	3.12690	1.05472	H17	-5.60791	1.95105	5.46082
F7	4.50842	1.20766	1.67932	C18	-0.31373	-9.37506	-0.87164	H18	-3.29294	1.64012	2.23622
F8	3.02616	-1.12710	3.49594	C19	-2.37995	5.40795	1.98170	H19	-5.12456	2.55028	3.28229
F9	1.91645	-2.82305	3.91156	C20	1.50659	-7.84023	-1.49625	H20	4.18054	-2.45276	-2.47461
				C21	-1.45849	-0.00777	3.02544	H21	1.26068	5.14226	2.62966
				C22	-0.35047	-0.53917	3.70609	H22	2.36022	1.51858	-3.38058
				C23	-0.42679	6.17678	3.03128	H23	0.18093	-11.2830	-1.15651
				C24	-2.94810	0.13921	5.06504	H24	2.30934	-10.9353	-1.99827
				C25	1.83165	3.04211	1.12943	H25	-4.31616	0.42752	6.53725
				C26	-2.36582	1.20483	-1.93990	H26	0.35052	2.81315	-7.30568
				C27	0.20071	-8.04939	-0.98839	H27	2.64570	2.90926	-7.12122
				C28	-3.83564	1.58076	-1.91421	H28	3.63501	2.19979	-5.18318
				C29	2.55808	-1.63871	-1.60249	C39	3.36083	-2.65628	-2.08467
				C30	-0.00563	2.18565	-5.42369	C40	4.06182	1.98497	0.77452
				C31	2.95945	-3.95933	-1.99594	C41	0.34769	5.19596	2.46037
				C32	-2.39409	-2.44994	0.25965	C42	1.94189	1.79841	-4.16280
				C33	0.75755	-1.00556	3.04178	C43	0.49750	-10.4124	-1.24057
				C34	-1.76093	6.31246	2.78727	C44	1.79490	-10.2032	-1.74259
				C35	-4.87300	1.58076	5.02768	C45	-4.09286	0.68757	5.67331
				C36	-3.48232	1.40269	3.11417	C46	0.76320	2.57572	-6.50712
				C37	1.91645	-1.55109	3.89522	C47	2.12561	2.61247	-6.40905
				C38	-4.58186	1.94539	3.73294	C48	2.70789	2.20544	-5.25090

TABLE IX: Experimental geometry, used in calculations, for complex **7**. All coordinates are expressed in Å.

Atom	x	y	z	Atom	x	y	z	Atom	x	y	z
Dy	0.00000	0.00000	0.00000	C1	6.17715	1.24526	2.25054	H1	7.07424	1.33300	1.94809
O1	1.98459	0.62446	1.06400	C2	5.80861	1.40253	3.54282	H2	6.43980	1.61773	4.21963
O2	-0.56902	0.51189	2.16594	C3	4.40941	1.22539	3.84104	H3	3.98265	1.30320	4.68494
O3	0.80980	-1.92326	0.99632	C4	3.78096	0.89430	2.54030	H4	1.85630	0.43574	4.23232
O4	-1.84916	-1.39186	0.38466	C5	2.37049	0.66420	2.26534	H5	4.23608	-4.35512	4.08215
O5	0.89905	1.64422	-1.36785	C6	1.48694	0.49865	3.35881	H6	2.27653	-5.14643	5.21370
O6	-1.64743	1.62601	-0.35306	C7	0.12810	0.42084	3.22345	H7	0.17874	-4.33360	4.20694
N1	-0.88082	-0.65519	-2.27562	C8	-0.72216	0.23212	4.48190	H8	-1.34407	-3.59030	2.64605
N2	1.67286	-1.02435	-1.62842	C9	3.34893	-4.13660	3.82201	H9	3.15440	-1.17997	-0.32979
F1	-0.00306	0.05996	5.59927	C10	2.24776	-4.58026	4.45229	H10	4.83974	-1.62197	-1.84627
F2	-1.49668	1.30982	4.70609	C11	1.04413	-4.12336	3.87277	H11	4.32320	-1.81897	-4.10724
F3	-1.53553	-0.81246	4.35922	C12	1.27655	-3.34033	2.76872	H12	2.12233	-1.45643	-4.79039
F4	-3.57598	-4.04721	1.56104	C13	0.32823	-2.69801	1.88464	H13	0.06552	-1.19321	-5.31492
F5	-4.00530	-2.20966	2.47262	C14	-1.05279	-2.95130	2.00519	H14	-2.19613	-0.87371	-5.73581
F6	-4.22832	-2.42983	0.43161	C15	-2.00020	-2.31229	1.23532	H15	-3.64119	-0.48799	-3.95284
F7	-4.11509	2.26502	-1.39577	C16	-3.45275	-2.75927	1.43625	H16	-2.77757	-0.33734	-1.79974
F8	-3.39092	3.76485	-0.05061	C17	2.94031	-1.22798	-1.25406	H17	4.42865	3.66884	-4.45833
F9	-3.33755	4.10919	-2.14449	C18	3.94710	-1.50609	-2.14872	H18	2.71765	5.01141	-5.37837
S1	4.88875	0.88007	1.23469	C19	3.64735	-1.60873	-3.47061	H19	0.34139	4.59092	-4.46468
S2	2.95824	-3.13091	2.49842	C20	2.34827	-1.40511	-3.86824	H20	-1.12535	3.64897	-2.75150
S3	3.03241	2.58253	-3.00933	C21	1.37115	-1.12534	-2.93763				
				C22	-0.04871	-0.92172	-3.29084				
				C23	-0.53083	-1.00946	-4.60004				
				C24	-1.86222	-0.83232	-4.84750				
				C25	-2.70853	-0.59228	-3.80267				
				C26	-2.18799	-0.50620	-2.52519				
				C27	3.52524	3.60924	-4.17280				
				C28	2.53233	4.37240	-4.70156				
				C29	1.15595	4.18534	-4.19395				
				C30	1.43854	3.12088	-3.08356				
				C31	0.47745	2.53154	-2.14660				
				C32	-0.85792	2.99341	-2.11911				
				C33	-1.77757	2.53982	-1.22234				
				C34	-3.15592	3.17386	-1.21811				

TABLE X: Experimental geometry, used in calculations, for complex **8**. All coordinates are expressed in Å.

Atom	x	y	z	Atom	x	y	z	Atom	x	y	z
Dy	0.00000	0.00000	0.00000	C1	-3.42833	-2.57014	-5.02550	H1	-4.32653	-2.83832	-5.18116
O1	-0.79991	-0.88383	-2.00300	C2	-2.44798	-2.64218	-5.96550	H2	-2.61998	-2.98610	-6.83447
O2	1.73433	0.06792	-1.53754	C3	-1.12534	-2.17027	-5.58618	H3	-0.33897	-2.12107	-6.11662
O3	-0.90087	-2.07916	0.47392	C4	-1.31864	-1.76578	-4.09457	H4	1.26068	-1.24447	-4.42251
O4	1.62800	-1.19372	1.19328	C5	-0.37032	-1.18783	-3.13644	H5	-4.22264	-6.06726	0.35151
O5	-0.81348	0.29703	2.14083	C6	0.98132	-0.95786	-3.55959	H6	-2.83365	-6.95021	2.13479
O6	1.19085	1.74699	0.95752	C7	1.86538	-0.33419	-2.73747	H7	-0.97407	-5.46102	2.71510
N1	-0.57641	2.16017	-1.25343	C8	3.25956	-0.00288	-3.31023	H8	0.52481	-3.70126	2.82240
N2	-2.47370	0.59592	-0.11848	C9	-3.46084	-5.62364	0.70514	H9	-3.85406	-0.77315	6.08972
F1	4.23090	-0.53023	-2.55914	C10	-2.67614	-6.11608	1.70710	H10	-2.67614	0.83163	7.32139
F2	3.44417	-0.43821	-4.54039	C11	-1.61432	-5.26596	2.04109	H11	-0.80337	2.17505	6.05950
F3	3.45525	1.30342	-3.30721	C12	-1.60081	-4.12441	1.29000	H12	0.36111	2.66420	3.89539
F4	2.67639	-3.35505	3.72613	C13	-0.65236	-3.01112	1.28849	H13	-3.15016	-0.97776	0.85476
F5	3.76376	-2.89922	1.93681	C14	0.48498	-3.04993	2.13176	H14	-5.40921	-0.38572	0.83511
F6	3.28686	-1.35503	3.32263	C15	1.52960	-2.17693	1.98064	H15	-6.08524	1.61204	-0.06407
F7	1.91528	4.31657	3.19568	C16	2.80666	-2.44519	2.77555	H16	-5.42367	3.76638	-1.23227
F8	3.35717	3.10334	2.16501	C17	-3.15626	-0.25077	5.70889	H17	-3.83479	5.08941	-2.15111
F9	1.97810	4.30513	1.03762	C18	-2.47095	0.67911	6.40708	H18	-1.39211	5.44590	-2.80094
S1	-2.89242	-1.96967	-3.59586	C19	-1.36964	1.48581	5.73307	H19	0.78086	4.68732	-2.77676
S2	-2.89924	-4.11474	0.16986	C20	-1.47417	0.80992	4.33063	H20	1.26512	2.63446	-1.78992
S3	-2.67407	-0.40198	4.13266	C21	-0.64379	1.05540	3.12768				
				C22	0.29624	2.10842	3.12919				
				C23	1.12134	2.35458	2.05469				
				C24	2.08013	3.54129	2.12874				
				C25	-3.40861	-0.16101	0.44370				
				C26	-4.76888	0.20204	0.45125				
				C27	-5.16823	1.36293	-0.07314				
				C28	-4.20177	2.20324	-0.63986				
				C29	-2.86775	1.76394	-0.65950				
				C30	-4.52037	3.47080	-1.23227				
				C31	-3.58457	4.24780	-1.78538				
				C32	-2.20992	3.84076	-1.83979				
				C33	-1.85994	2.60499	-1.27156				
				C34	-1.18791	4.60673	-2.40349				
				C35	0.08516	4.16729	-2.39140				
				C36	0.36495	2.93698	-1.80201				

TABLE XI: Experimental geometry, used in calculations, for complex **9**. All coordinates are expressed in Å.

Atom	x	y	z	Atom	x	y	z	Atom	x	y	z
Dy	0.00000	0.00000	0.00000	C1	3.06851	0.79179	1.03378	H1	5.75045	0.94581	2.08534
O1	2.10599	0.98194	0.21859	C2	4.12948	1.81099	1.12325	H2	6.87864	3.15155	1.91965
O2	1.38353	-1.61931	0.98186	C3	5.39117	1.70831	1.68880	H3	5.59188	4.76592	0.61735
O3	0.54858	0.83172	-2.12865	C4	6.04103	2.96901	1.55845	H4	3.79612	-0.40045	2.57909
O4	-0.12979	-1.80756	-1.54764	C5	5.28637	3.91215	0.82612	H5	-0.88624	1.01426	-6.34374
O5	0.10345	1.04089	2.12952	C6	3.16867	-0.38524	1.89314	H6	-0.52409	3.39684	-6.85737
O6	-1.51369	-1.12873	1.43474	C7	2.38627	-1.41775	1.71089	H7	0.65440	4.80205	-5.28776
N1	-0.87477	2.39856	-0.25527	C8	2.62306	-2.67655	2.60229	H8	-0.57582	-0.88724	-4.51013
N2	-2.33590	0.29169	-1.00087	C9	0.15499	0.53128	-3.29399	H9	-2.06400	2.39475	5.23120
F1	3.46345	-2.46168	3.56659	C10	0.05573	1.61134	-4.38531	H10	-0.79278	3.93307	6.68815
F2	3.07353	-3.63110	1.90088	C11	-0.50511	1.66838	-5.81023	H11	1.33311	4.36471	5.84866
F3	1.52152	-3.10438	3.17004	C12	-0.24945	3.03746	-6.04219	H12	-2.31846	0.41339	4.03935
F4	-2.60012	-2.95987	3.25841	C13	0.41958	3.91215	-5.11434	H13	0.68824	3.20860	0.68583
F5	-3.87412	-1.42726	3.92558	C14	-0.26863	-0.75223	-3.64193	H14	0.19397	5.39722	-0.01779
F6	-3.93416	-2.13272	1.93401	C15	-0.26287	-1.79235	-2.81791	H15	-1.63816	5.86689	-1.21626
F7	0.17358	-4.05133	-2.94273	C16	-0.59487	-3.18044	-3.36799	H16	-3.05496	4.12512	-1.87791
F8	-0.69280	-3.25270	-4.66698	C17	-0.65892	1.07131	3.16452	H17	-4.30176	2.36813	-2.35288
F9	-1.73261	-3.64441	-2.91401	C18	-0.34548	2.10192	4.09679	H18	-2.67057	-1.59459	-0.99534
S1	3.81752	3.34246	0.38539	C19	-1.16940	2.61913	5.10969	H19	-6.74513	1.04279	-2.47107
S2	0.69428	3.12303	-3.73361	C20	-0.44791	3.53185	5.92377	H20	-5.94723	0.95722	-3.82308
S3	1.19595	2.77981	4.12772	C21	0.75709	3.73911	5.46979	H21	-7.45395	-0.87583	-3.65408
				C22	-1.74692	0.23845	3.32680	H22	-4.71227	-2.80205	-1.92099
				C23	-2.04663	-0.80927	2.53380	H23	-7.14967	-2.42935	-1.87791
				C24	-3.13980	-1.79805	2.91709	H24	-6.71625	-1.11161	-1.07377
				C25	-0.05462	3.38353	0.15453	H25	-3.89635	-2.01483	-4.97848
				C26	-0.37452	4.70317	-0.26190	H26	-4.04085	-0.56208	-4.40409
				C27	-1.44844	4.98650	-0.98209	H27	-4.97904	-1.04316	-5.56391
				C28	-2.28523	3.95969	-1.38195	H28	-6.70371	-2.99980	-4.93761
				C29	-1.92015	2.65336	-1.00308	H29	-6.51227	-3.62159	-3.51159
				C30	-2.82682	1.49725	-1.35102	H30	-5.34755	-3.68815	-4.55763
				C31	-4.01588	1.55049	-2.01377				
				C32	-4.79937	0.45522	-2.20045	C37	-5.06857	-1.93496	-2.20928
				C33	-4.26588	-0.75413	-1.89116	C38	-5.54386	-1.82848	-3.72146
				C34	-3.02223	-0.77315	-1.25271	C39	-6.58040	-1.64403	-1.87238
				C35	-6.09275	0.52368	-2.96813	C40	-4.51721	-1.31317	-4.76640
				C36	-6.59746	-0.84160	-3.17911	C41	-6.07635	-3.15573	-4.22846

TABLE XII: Experimental geometry, used in calculations, for complex **10**. All coordinates are expressed in Å.

COMPUTATIONAL RESULTS

Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|
O1	$-5.24\cdot10^{6}$	$1.29\cdot10^{-15}$	2.325	C1	$-5.13\cdot10^{5}$	$3.72\cdot10^{-17}$	0.067	H1	$-6.26\cdot10^{5}$	$7.16\cdot10^{-16}$	1.295
O2	$8.52\cdot10^{6}$	$2.09\cdot10^{-15}$	3.782	C2	$-2.94\cdot10^{5}$	$2.13\cdot10^{-17}$	0.039	H2	$-1.54\cdot10^{5}$	$1.76\cdot10^{-16}$	0.318
O3	$-2.63\cdot10^{5}$	$6.46\cdot10^{-17}$	0.117	C3	$1.30\cdot10^{6}$	$9.42\cdot10^{-17}$	0.170	H3	$-6.27\cdot10^{5}$	$7.19\cdot10^{-16}$	1.298
O4	$-2.11\cdot10^{4}$	$5.17\cdot10^{-18}$	0.009	C4	$3.00\cdot10^{6}$	$2.17\cdot10^{-16}$	0.392	H4	$6.84\cdot10^{5}$	$7.84\cdot10^{-16}$	1.416
O5	$6.30\cdot10^{6}$	$1.55\cdot10^{-15}$	2.795	C5	$1.12\cdot10^{6}$	$8.09\cdot10^{-17}$	0.146	H5	$8.19\cdot10^{5}$	$9.38\cdot10^{-16}$	1.694
O6	$-8.19\cdot10^{6}$	$2.01\cdot10^{-15}$	3.634	C6	$4.26\cdot10^{5}$	$3.09\cdot10^{-17}$	0.056	H6	$7.65\cdot10^{5}$	$8.76\cdot10^{-16}$	1.583
O7	$1.11\cdot10^{5}$	$2.72\cdot10^{-17}$	0.049	C7	$9.29\cdot10^{5}$	$6.72\cdot10^{-17}$	0.122	H7	$9.49\cdot10^{5}$	$1.09\cdot10^{-15}$	1.965
O8	$1.55\cdot10^{5}$	$3.81\cdot10^{-17}$	0.069	C8	$6.82\cdot10^{5}$	$4.94\cdot10^{-17}$	0.089	H8	$3.20\cdot10^{4}$	$3.66\cdot10^{-17}$	0.066
				C9	$3.11\cdot10^{5}$	$2.25\cdot10^{-17}$	0.041	H9	$3.76\cdot10^{5}$	$4.30\cdot10^{-16}$	0.778
				C10	$2.29\cdot10^{4}$	$1.66\cdot10^{-18}$	0.003	H10	$6.37\cdot10^{5}$	$7.29\cdot10^{-16}$	1.317
				C11	$7.94\cdot10^{5}$	$5.74\cdot10^{-17}$	0.104	H11	$4.18\cdot10^{5}$	$4.79\cdot10^{-16}$	0.865
				C12	$1.06\cdot10^{6}$	$7.66\cdot10^{-17}$	0.138	H12	$-9.76\cdot10^{4}$	$1.12\cdot10^{-16}$	0.202
				C13	$-7.99\cdot10^{5}$	$5.78\cdot10^{-17}$	0.105	H13	$7.39\cdot10^{4}$	$8.46\cdot10^{-17}$	0.153
				C14	$-2.64\cdot10^{6}$	$1.91\cdot10^{-16}$	0.345	H14	$2.30\cdot10^{3}$	$2.63\cdot10^{-18}$	0.005
				C15	$-1.02\cdot10^{6}$	$7.37\cdot10^{-17}$	0.133	H15	$1.02\cdot10^{6}$	$1.17\cdot10^{-15}$	2.107
								H16	$3.89\cdot10^{5}$	$4.45\cdot10^{-16}$	0.805
								H17	$6.33\cdot10^{5}$	$7.25\cdot10^{-16}$	1.310
								H18	$-3.73\cdot10^{5}$	$4.27\cdot10^{-16}$	0.772
								H19	$-8.85\cdot10^{5}$	$1.01\cdot10^{-15}$	1.832
								H20	$-6.25\cdot10^{5}$	$7.16\cdot10^{-16}$	1.294
								H21	$-8.15\cdot10^{5}$	$9.33\cdot10^{-16}$	1.686
								H22	$-7.70\cdot10^{5}$	$8.82\cdot10^{-16}$	1.594
								H23	$4.95\cdot10^{5}$	$5.67\cdot10^{-16}$	1.024
								H24	$1.12\cdot10^{6}$	$1.28\cdot10^{-15}$	2.310
								H25	$-2.01\cdot10^{5}$	$2.31\cdot10^{-16}$	0.417

TABLE XIII: Values of $\beta^2\sigma^{(1p)}$, expressed in ppm au, of the rotating electric polarization, expressed in $\frac{C}{m^2}$, and of the induced voltage, expressed in μV, for non-lanthanide atoms of complex **1**. 1 ppm au of $\beta^2\sigma^{(1p)}$ $= 1.94469097 \cdot 10^{-18}\frac{m}{V}$.

Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|
O1	$-2.11\cdot10^{4}$	$5.18\cdot10^{-18}$	0.009	C1	$1.92\cdot10^{5}$	$1.39\cdot10^{-17}$	0.025	H1	$1.49\cdot10^{4}$	$1.71\cdot10^{-17}$	0.031
O2	$9.43\cdot10^{6}$	$2.32\cdot10^{-15}$	4.183	C2	$8.06\cdot10^{3}$	$5.83\cdot10^{-19}$	0.001	H2	$-2.65\cdot10^{3}$	$3.04\cdot10^{-18}$	0.005
O3	$-6.06\cdot10^{6}$	$1.49\cdot10^{-15}$	2.687	C3	$5.95\cdot10^{3}$	$4.31\cdot10^{-19}$	0.001	H3	$1.11\cdot10^{6}$	$1.27\cdot10^{-15}$	2.292
O4	$7.34\cdot10^{6}$	$1.80\cdot10^{-15}$	3.258	C4	$1.32\cdot10^{5}$	$9.58\cdot10^{-18}$	0.017	H4	$3.85\cdot10^{5}$	$4.41\cdot10^{-16}$	0.796
O5	$-1.35\cdot10^{6}$	$3.31\cdot10^{-16}$	0.599	C5	$-2.59\cdot10^{3}$	$1.88\cdot10^{-19}$	<0.001	H5	$6.78\cdot10^{5}$	$7.77\cdot10^{-16}$	1.403
O6	$-1.02\cdot10^{7}$	$2.50\cdot10^{-15}$	4.513	C6	$8.34\cdot10^{5}$	$6.04\cdot10^{-17}$	0.109	H6	$-5.96\cdot10^{3}$	$6.82\cdot10^{-18}$	0.012
N1	$1.48\cdot10^{6}$	$2.36\cdot10^{-17}$	0.043	C7	$-7.64\cdot10^{5}$	$5.53\cdot10^{-17}$	0.100	H7	$2.88\cdot10^{4}$	$3.30\cdot10^{-17}$	0.060
N2	$7.94\cdot10^{5}$	$1.27\cdot10^{-17}$	0.023	C8	$3.10\cdot10^{6}$	$2.24\cdot10^{-16}$	0.405	H8	$1.26\cdot10^{5}$	$1.44\cdot10^{-16}$	0.261
N3	$2.63\cdot10^{4}$	$4.19\cdot10^{-19}$	0.001	C9	$4.81\cdot10^{5}$	$3.48\cdot10^{-17}$	0.063	H9	$4.97\cdot10^{5}$	$5.69\cdot10^{-16}$	1.028
N4	$-1.35\cdot10^{4}$	$2.16\cdot10^{-19}$	<0.001	C10	$-5.49\cdot10^{3}$	$3.97\cdot10^{-19}$	0.001	H10	$2.18\cdot10^{5}$	$2.49\cdot10^{-16}$	0.450
				C11	$-1.71\cdot10^{4}$	$1.24\cdot10^{-18}$	0.002	H11	$1.65\cdot10^{6}$	$1.89\cdot10^{-15}$	3.418
				C12	$-2.40\cdot10^{4}$	$1.74\cdot10^{-18}$	0.003	H12	$2.81\cdot10^{3}$	$3.22\cdot10^{-18}$	0.006
				C13	$4.84\cdot10^{4}$	$3.50\cdot10^{-18}$	0.006	H13	$3.47\cdot10^{5}$	$3.97\cdot10^{-16}$	0.718
				C14	$1.99\cdot10^{5}$	$1.44\cdot10^{-17}$	0.026	H14	$3.04\cdot10^{5}$	$3.48\cdot10^{-16}$	0.629
				C15	$-3.38\cdot10^{6}$	$2.45\cdot10^{-16}$	0.442	H15	$-1.92\cdot10^{4}$	$2.20\cdot10^{-17}$	0.040
				C16	$2.40\cdot10^{4}$	$1.74\cdot10^{-18}$	0.003	H16	$5.63\cdot10^{4}$	$6.45\cdot10^{-17}$	0.117
				C17	$-5.08\cdot10^{4}$	$3.67\cdot10^{-18}$	0.007	H17	$-6.48\cdot10^{3}$	$7.42\cdot10^{-18}$	0.013
				C18	$1.01\cdot10^{6}$	$7.30\cdot10^{-17}$	0.132	H18	$-8.10\cdot10^{5}$	$9.28\cdot10^{-16}$	1.677
				C19	$-6.50\cdot10^{3}$	$4.71\cdot10^{-19}$	0.001	H19	$-2.29\cdot10^{5}$	$2.62\cdot10^{-16}$	0.473
				C20	$2.35\cdot10^{5}$	$1.70\cdot10^{-17}$	0.031	H20	$-7.27\cdot10^{5}$	$8.32\cdot10^{-16}$	1.504
				C21	$1.02\cdot10^{6}$	$7.37\cdot10^{-17}$	0.133	H21	$8.45\cdot10^{5}$	$9.67\cdot10^{-16}$	1.748
				C22	$2.67\cdot10^{5}$	$1.93\cdot10^{-17}$	0.035	H22	$7.06\cdot10^{5}$	$8.08\cdot10^{-16}$	1.460
				C23	$9.10\cdot10^{5}$	$6.59\cdot10^{-17}$	0.119	H23	$1.01\cdot10^{6}$	$1.15\cdot10^{-15}$	2.084
				C24	$1.92\cdot10^{3}$	$1.39\cdot10^{-19}$	<0.001	H24	$1.26\cdot10^{6}$	$1.44\cdot10^{-15}$	2.606
				C25	$5.23\cdot10^{5}$	$3.79\cdot10^{-17}$	0.068	H25	$-1.11\cdot10^{6}$	$1.27\cdot10^{-15}$	2.290
				C26	$3.96\cdot10^{5}$	$2.87\cdot10^{-17}$	0.052	H26	$-8.88\cdot10^{5}$	$1.02\cdot10^{-15}$	1.838
				C27	$2.62\cdot10^{4}$	$1.89\cdot10^{-18}$	0.003	H27	$-8.79\cdot10^{5}$	$1.01\cdot10^{-15}$	1.820
				C28	$-6.75\cdot10^{5}$	$4.89\cdot10^{-17}$	0.088	H28	$-5.90\cdot10^{5}$	$6.76\cdot10^{-16}$	1.221
				C29	$1.13\cdot10^{6}$	$8.19\cdot10^{-17}$	0.148	H29	$-1.58\cdot10^{5}$	$1.80\cdot10^{-16}$	0.326
				C30	$1.06\cdot10^{6}$	$7.64\cdot10^{-17}$	0.138	H30	$1.29\cdot10^{5}$	$1.48\cdot10^{-16}$	0.268
				C31	$-1.26\cdot10^{6}$	$9.15\cdot10^{-17}$	0.165	H31	$5.59\cdot10^{5}$	$6.40\cdot10^{-16}$	1.157
				C32	$-1.20\cdot10^{6}$	$8.68\cdot10^{-17}$	0.157				
				C33	$2.23\cdot10^{5}$	$1.61\cdot10^{-17}$	0.029				

TABLE XIV: Values of $\beta^2\sigma^{(1p)}$, expressed in ppm au, of the rotating electric polarization, expressed in $\frac{C}{m^2}$, and of the induced voltage, expressed in μV, for non-lanthanide atoms of complex **2**. 1 ppm au of $\beta^2\sigma^{(1p)}$ = $1.94469097\cdot10^{-18}\frac{m}{V}$.

Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|
O1	$2.16\cdot10^{6}$	$5.31\cdot10^{-16}$	0.959	C1	$-4.55\cdot10^{5}$	$3.29\cdot10^{-17}$	0.059	H1	$-4.39\cdot10^{5}$	$5.03\cdot10^{-16}$	0.909
O2	$-9.22\cdot10^{6}$	$2.26\cdot10^{-15}$	4.091	C2	$-1.86\cdot10^{5}$	$1.34\cdot10^{-17}$	0.024	H2	$-1.45\cdot10^{5}$	$1.67\cdot10^{-16}$	0.301
O3	$3.98\cdot10^{4}$	$9.76\cdot10^{-18}$	0.018	C3	$-9.78\cdot10^{4}$	$7.08\cdot10^{-18}$	0.013	H3	$-6.42\cdot10^{4}$	$7.36\cdot10^{-17}$	0.133
O4	$5.56\cdot10^{6}$	$1.36\cdot10^{-15}$	2.466	C4	$-6.00\cdot10^{4}$	$4.34\cdot10^{-18}$	0.008	H4	$-2.17\cdot10^{3}$	$2.49\cdot10^{-18}$	0.004
O5	$-9.70\cdot10^{6}$	$2.38\cdot10^{-15}$	4.303	C5	$-9.71\cdot10^{3}$	$7.03\cdot10^{-19}$	0.001	H5	$1.55\cdot10^{3}$	$1.78\cdot10^{-18}$	0.003
O6	$1.21\cdot10^{7}$	$2.96\cdot10^{-15}$	5.350	C6	$-1.97\cdot10^{3}$	$1.43\cdot10^{-19}$	<0.001	H6	$2.33\cdot10^{4}$	$2.67\cdot10^{-17}$	0.048
N1	$-7.10\cdot10^{5}$	$1.13\cdot10^{-17}$	0.020	C7	$1.83\cdot10^{3}$	$1.32\cdot10^{-19}$	<0.001	H7	$3.88\cdot10^{4}$	$4.45\cdot10^{-17}$	0.080
N2	$1.72\cdot10^{5}$	$2.74\cdot10^{-18}$	0.005	C8	$8.84\cdot10^{3}$	$6.40\cdot10^{-19}$	0.001	H8	$-1.64\cdot10^{5}$	$1.88\cdot10^{-16}$	0.339
N3	$-8.96\cdot10^{3}$	$1.43\cdot10^{-19}$	<0.001	C9	$3.78\cdot10^{4}$	$2.74\cdot10^{-18}$	0.005	H9	$-1.29\cdot10^{6}$	$1.48\cdot10^{-15}$	2.666
N4	$5.98\cdot10^{3}$	$9.54\cdot10^{-20}$	<0.001	C10	$3.80\cdot10^{4}$	$2.75\cdot10^{-18}$	0.005	H10	$-5.97\cdot10^{5}$	$6.83\cdot10^{-16}$	1.235
811				C11	$5.01\cdot10^{4}$	$3.63\cdot10^{-18}$	0.007	H11	$-8.70\cdot10^{5}$	$9.96\cdot10^{-16}$	1.800
				C12	$2.94\cdot10^{4}$	$2.13\cdot10^{-18}$	0.004	H12	$3.45\cdot10^{5}$	$3.95\cdot10^{-16}$	0.713
				C13	$7.69\cdot10^{4}$	$5.56\cdot10^{-18}$	0.010	H13	$1.25\cdot10^{6}$	$1.43\cdot10^{-15}$	2.585
				C14	$-1.06\cdot10^{5}$	$7.68\cdot10^{-18}$	0.014	H14	$8.55\cdot10^{5}$	$9.80\cdot10^{-16}$	1.770
				C15	$-1.11\cdot10^{6}$	$8.02\cdot10^{-17}$	0.145	H15	$1.29\cdot10^{6}$	$1.47\cdot10^{-15}$	2.660
				C16	$-1.94\cdot10^{6}$	$1.40\cdot10^{-16}$	0.253	H16	$-9.41\cdot10^{5}$	$1.08\cdot10^{-15}$	1.948
				C17	$6.50\cdot10^{5}$	$4.70\cdot10^{-17}$	0.085	H17	$-9.20\cdot10^{5}$	$1.05\cdot10^{-15}$	1.904
				C18	$3.33\cdot10^{6}$	$2.41\cdot10^{-16}$	0.436	H18	$-9.11\cdot10^{5}$	$1.04\cdot10^{-15}$	1.885
				C19	$1.45\cdot10^{6}$	$1.05\cdot10^{-16}$	0.189	H19	$-1.25\cdot10^{6}$	$1.43\cdot10^{-15}$	2.591
				C20	$-1.28\cdot10^{6}$	$9.24\cdot10^{-17}$	0.167	H20	$3.55\cdot10^{5}$	$4.07\cdot10^{-16}$	0.735
				C21	$-3.90\cdot10^{6}$	$2.82\cdot10^{-16}$	0.510	H21	$4.04\cdot10^{5}$	$4.63\cdot10^{-16}$	0.836
				C22	$-2.43\cdot10^{6}$	$1.76\cdot10^{-16}$	0.318	H22	$-2.37\cdot10^{5}$	$2.72\cdot10^{-16}$	0.491
				C23	$-1.31\cdot10^{6}$	$9.51\cdot10^{-17}$	0.172	H23	$7.06\cdot10^{4}$	$8.09\cdot10^{-17}$	0.146
				C24	$5.49\cdot10^{4}$	$3.98\cdot10^{-18}$	0.007	H24	$5.61\cdot10^{5}$	$6.42\cdot10^{-16}$	1.161
				C25	$6.02\cdot10^{5}$	$4.36\cdot10^{-17}$	0.079	H25	$3.85\cdot10^{5}$	$4.41\cdot10^{-16}$	0.798
				C26	$1.70\cdot10^{6}$	$1.23\cdot10^{-16}$	0.223	H26	$9.64\cdot10^{5}$	$1.10\cdot10^{-15}$	1.996
				C27	$-1.54\cdot10^{4}$	$1.11\cdot10^{-18}$	0.002	H27	$7.73\cdot10^{3}$	$8.85\cdot10^{-18}$	0.016
				C28	$8.00\cdot10^{5}$	$5.79\cdot10^{-17}$	0.105	H28	$-3.25\cdot10^{4}$	$3.72\cdot10^{-17}$	0.067
				C29	$-5.77\cdot10^{2}$	$4.18\cdot10^{-20}$	<0.001	H29	$3.89\cdot10^{4}$	$4.45\cdot10^{-17}$	0.081

TABLE XV: Values of $\beta^2\sigma^{(1p)}$, expressed in ppm au, of the rotating electric polarization, expressed in $\frac{C}{m^2}$, and of the induced voltage, expressed in μV, for non-lanthanide atoms of complex **3**. 1 ppm au of $\beta^2\sigma^{(1p)}$ = $1.94469097\cdot10^{-18}\frac{m}{V}$.

Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|
O1	$-1.03\cdot10^{7}$	$2.53\cdot10^{-15}$	4.565	C1	$-8.61\cdot10^{5}$	$6.23\cdot10^{-17}$	0.113	H1	$-1.65\cdot10^{6}$	$1.89\cdot10^{-15}$	3.413
O2	$1.36\cdot10^{7}$	$3.34\cdot10^{-15}$	6.030	C2	$-1.82\cdot10^{5}$	$1.32\cdot10^{-17}$	0.024	H2	$-1.79\cdot10^{5}$	$2.05\cdot10^{-16}$	0.371
O3	$4.48\cdot10^{6}$	$1.10\cdot10^{-15}$	1.988	C3	$-8.56\cdot10^{3}$	$6.20\cdot10^{-19}$	0.001	H3	$3.26\cdot10^{3}$	$3.74\cdot10^{-18}$	0.007
O4	$-1.46\cdot10^{3}$	$3.58\cdot10^{-19}$	0.001	C4	$4.31\cdot10^{4}$	$3.12\cdot10^{-18}$	0.006	H4	$1.83\cdot10^{4}$	$2.10\cdot10^{-17}$	0.038
O5	$5.20\cdot10^{6}$	$1.28\cdot10^{-15}$	2.309	C5	$1.87\cdot10^{4}$	$1.35\cdot10^{-18}$	0.002	H5	$-2.72\cdot10^{4}$	$3.11\cdot10^{-17}$	0.056
O6	$-1.17\cdot10^{7}$	$2.88\cdot10^{-15}$	5.200	C6	$-2.61\cdot10^{4}$	$1.89\cdot10^{-18}$	0.003	H6	$-1.21\cdot10^{5}$	$1.39\cdot10^{-16}$	0.251
N1	$-7.73\cdot10^{5}$	$1.23\cdot10^{-17}$	0.022	C7	$-1.33\cdot10^{5}$	$9.61\cdot10^{-18}$	0.017	H7	$-2.33\cdot10^{5}$	$2.67\cdot10^{-16}$	0.482
N2	$-1.24\cdot10^{6}$	$1.98\cdot10^{-17}$	0.036	C8	$-1.87\cdot10^{5}$	$1.36\cdot10^{-17}$	0.025	H8	$-8.31\cdot10^{5}$	$9.52\cdot10^{-16}$	1.720
				C9	$-3.21\cdot10^{5}$	$2.32\cdot10^{-17}$	0.042	H9	$-8.00\cdot10^{5}$	$9.16\cdot10^{-16}$	1.656
				C10	$-8.08\cdot10^{5}$	$5.85\cdot10^{-17}$	0.106	H10	$-5.82\cdot10^{5}$	$6.67\cdot10^{-16}$	1.205
				C11	$-2.13\cdot10^{5}$	$1.54\cdot10^{-17}$	0.028	H11	$-1.47\cdot10^{6}$	$1.69\cdot10^{-15}$	3.049
				C12	$7.01\cdot10^{4}$	$5.07\cdot10^{-18}$	0.009	H12	$6.55\cdot10^{5}$	$7.50\cdot10^{-16}$	1.356
				C13	$-1.11\cdot10^{6}$	$8.04\cdot10^{-17}$	0.145	H13	$1.45\cdot10^{6}$	$1.67\cdot10^{-15}$	3.011
				C14	$-1.55\cdot10^{6}$	$1.12\cdot10^{-16}$	0.203	H14	$1.05\cdot10^{6}$	$1.20\cdot10^{-15}$	2.177
				C15	$1.28\cdot10^{6}$	$9.24\cdot10^{-17}$	0.167	H15	$1.37\cdot10^{6}$	$1.57\cdot10^{-15}$	2.844
				C16	$4.14\cdot10^{6}$	$3.00\cdot10^{-16}$	0.542	H16	$-4.96\cdot10^{5}$	$5.68\cdot10^{-16}$	1.026
				C17	$1.68\cdot10^{6}$	$1.21\cdot10^{-16}$	0.220	H17	$2.89\cdot10^{4}$	$3.31\cdot10^{-17}$	0.060
				C18	$-2.51\cdot10^{4}$	$1.82\cdot10^{-18}$	0.003	H18	$4.56\cdot10^{5}$	$5.22\cdot10^{-16}$	0.944
				C19	$-1.94\cdot10^{5}$	$1.41\cdot10^{-17}$	0.025	H19	$-4.46\cdot10^{5}$	$5.11\cdot10^{-16}$	0.923
				C20	$-7.19\cdot10^{5}$	$5.20\cdot10^{-17}$	0.094	H20	$-2.35\cdot10^{3}$	$2.69\cdot10^{-18}$	0.005
				C21	$-6.06\cdot10^{5}$	$4.39\cdot10^{-17}$	0.079	H21	$-1.64\cdot10^{5}$	$1.87\cdot10^{-16}$	0.339
				C22	$-9.50\cdot10^{4}$	$6.88\cdot10^{-18}$	0.012	H22	$5.68\cdot10^{4}$	$6.50\cdot10^{-17}$	0.117
				C23	$2.68\cdot10^{5}$	$1.94\cdot10^{-17}$	0.035	H23	$5.62\cdot10^{5}$	$6.44\cdot10^{-16}$	1.164
				C24	$-4.73\cdot10^{5}$	$3.43\cdot10^{-17}$	0.062	H24	$-1.06\cdot10^{5}$	$1.21\cdot10^{-16}$	0.218
				C25	$-2.25\cdot10^{6}$	$1.63\cdot10^{-16}$	0.294	H25	$4.11\cdot10^{5}$	$4.70\cdot10^{-16}$	0.850
				C26	$-4.26\cdot10^{6}$	$3.09\cdot10^{-16}$	0.558	H26	$-1.19\cdot10^{6}$	$1.36\cdot10^{-15}$	2.456
				C27	$-1.54\cdot10^{6}$	$1.11\cdot10^{-16}$	0.201	H27	$-1.25\cdot10^{6}$	$1.43\cdot10^{-15}$	2.586
								H28	$-1.02\cdot10^{6}$	$1.17\cdot10^{-15}$	2.119
								H29	$-1.19\cdot10^{6}$	$1.36\cdot10^{-15}$	2.466

TABLE XVI: Values of $\beta^2\sigma^{(1p)}$, expressed in ppm au, of the rotating electric polarization, expressed in $\frac{C}{m^2}$, and of the induced voltage, expressed in μV, for non-lanthanide atoms of complex **4**. 1 ppm au of $\beta^2\sigma^{(1p)}$ = $1.94469097\cdot10^{-18}\frac{m}{V}$.

Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|
O1	$-6.45\cdot10^5$	$1.58\cdot10^{-16}$	0.286	C1	$5.41\cdot10^4$	$3.92\cdot10^{-18}$	0.007	H1	$2.27\cdot10^5$	$2.60\cdot10^{-16}$	0.469
O2	$1.46\cdot10^6$	$3.59\cdot10^{-16}$	0.648	C2	$-1.30\cdot10^6$	$9.44\cdot10^{-17}$	0.171	H2	$3.07\cdot10^4$	$3.52\cdot10^{-17}$	0.064
O3	$5.70\cdot10^6$	$1.40\cdot10^{-15}$	2.527	C3	$1.34\cdot10^4$	$9.67\cdot10^{-19}$	0.002	H3	$-1.81\cdot10^4$	$2.07\cdot10^{-17}$	0.037
O4	$-8.47\cdot10^6$	$2.08\cdot10^{-15}$	3.758	C4	$1.97\cdot10^6$	$1.42\cdot10^{-16}$	0.257	H4	$-2.29\cdot10^6$	$2.63\cdot10^{-15}$	4.748
O5	$-4.20\cdot10^6$	$1.03\cdot10^{-15}$	1.865	C5	$2.41\cdot10^5$	$1.74\cdot10^{-17}$	0.031	H5	$-4.24\cdot10^5$	$4.86\cdot10^{-16}$	0.878
O6	$9.79\cdot10^6$	$2.40\cdot10^{-15}$	4.342	C6	$-1.37\cdot10^6$	$9.92\cdot10^{-17}$	0.179	H6	$-4.14\cdot10^5$	$4.74\cdot10^{-16}$	0.857
O7	$-2.32\cdot10^6$	$5.71\cdot10^{-16}$	1.031	C7	$-3.18\cdot10^6$	$2.30\cdot10^{-16}$	0.416	H7	$1.06\cdot10^6$	$1.21\cdot10^{-15}$	2.191
O8	$-6.36\cdot10^6$	$1.56\cdot10^{-15}$	2.819	C8	$4.74\cdot10^5$	$3.43\cdot10^{-17}$	0.062	H8	$2.43\cdot10^6$	$2.79\cdot10^{-15}$	5.037
F1	$2.70\cdot10^5$	$2.74\cdot10^{-16}$	0.495	C9	$-1.27\cdot10^6$	$9.18\cdot10^{-17}$	0.166	H9	$1.87\cdot10^6$	$2.14\cdot10^{-15}$	3.862
F2	$7.88\cdot10^4$	$8.00\cdot10^{-17}$	0.145	C10	$-1.22\cdot10^6$	$8.84\cdot10^{-17}$	0.160	H10	$5.33\cdot10^5$	$6.11\cdot10^{-16}$	1.103
F3	$8.76\cdot10^5$	$8.89\cdot10^{-16}$	1.607	C11	$2.25\cdot10^5$	$1.63\cdot10^{-17}$	0.029	H11	$-7.55\cdot10^5$	$8.65\cdot10^{-16}$	1.563
F4	$-6.90\cdot10^5$	$7.00\cdot10^{-16}$	1.265	C12	$2.34\cdot10^6$	$1.69\cdot10^{-16}$	0.306	H12	$1.95\cdot10^4$	$2.24\cdot10^{-17}$	0.040
F5	$-1.09\cdot10^6$	$1.11\cdot10^{-15}$	2.005	C13	$-7.91\cdot10^5$	$5.72\cdot10^{-17}$	0.103	H13	$-1.06\cdot10^6$	$1.21\cdot10^{-15}$	2.186
F6	$-6.67\cdot10^5$	$6.77\cdot10^{-16}$	1.223	C14	$9.87\cdot10^5$	$7.14\cdot10^{-17}$	0.129				
F7	$-1.09\cdot10^6$	$1.10\cdot10^{-15}$	1.993	C15	$-1.84\cdot10^6$	$1.33\cdot10^{-16}$	0.241				
F8	$-3.55\cdot10^5$	$3.60\cdot10^{-16}$	0.650	C16	$-2.02\cdot10^6$	$1.46\cdot10^{-16}$	0.264				
F9	$-3.10\cdot10^5$	$3.14\cdot10^{-16}$	0.568	C17	$-3.01\cdot10^6$	$2.18\cdot10^{-16}$	0.394				
F10	$7.20\cdot10^5$	$7.31\cdot10^{-16}$	1.320	C18	$-4.17\cdot10^5$	$3.02\cdot10^{-17}$	0.055				
F11	$9.78\cdot10^5$	$9.92\cdot10^{-16}$	1.793	C19	$-1.02\cdot10^6$	$7.39\cdot10^{-17}$	0.134				
F12	$3.62\cdot10^5$	$3.67\cdot10^{-16}$	0.663								
F13	$-2.67\cdot10^4$	$2.71\cdot10^{-17}$	0.049								
F14	$-1.62\cdot10^5$	$1.64\cdot10^{-16}$	0.297								
F15	$-3.58\cdot10^5$	$3.63\cdot10^{-16}$	0.656								
F16	$-2.20\cdot10^5$	$2.24\cdot10^{-16}$	0.404								
F17	$-1.01\cdot10^6$	$1.02\cdot10^{-15}$	1.848								
F18	$-6.52\cdot10^5$	$6.61\cdot10^{-16}$	1.195								

TABLE XVII: Values of $\beta^2\sigma^{(1p)}$, expressed in ppm au, of the rotating electric polarization, expressed in $\frac{C}{m^2}$, and of the induced voltage, expressed in μV, for non-lanthanide atoms of complex **5**. 1 ppm au of $\beta^2\sigma^{(1p)}$ $= 1.94469097 \cdot 10^{-18}\frac{m}{V}$.

Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|
O1	$2.74\cdot10^{7}$	$6.72\cdot10^{-15}$	12.15	C1	$6.04\cdot10^{4}$	$4.37\cdot10^{-18}$	0.008	H1	$1.32\cdot10^{6}$	$1.51\cdot10^{-15}$	2.734
O2	$1.92\cdot10^{7}$	$4.71\cdot10^{-15}$	8.504	C2	$-9.18\cdot10^{5}$	$6.65\cdot10^{-17}$	0.120	H2	$-1.46\cdot10^{6}$	$1.67\cdot10^{-15}$	3.018
O3	$-1.23\cdot10^{7}$	$3.03\cdot10^{-15}$	5.470	C3	$7.23\cdot10^{6}$	$5.23\cdot10^{-16}$	0.946	H3	$1.63\cdot10^{5}$	$1.87\cdot10^{-16}$	0.338
O4	$-2.46\cdot10^{6}$	$6.05\cdot10^{-16}$	1.093	C4	$5.03\cdot10^{6}$	$3.64\cdot10^{-16}$	0.658	H4	$-1.36\cdot10^{6}$	$1.56\cdot10^{-15}$	2.821
O5	$-2.21\cdot10^{6}$	$5.42\cdot10^{-16}$	0.980	C5	$6.96\cdot10^{5}$	$5.04\cdot10^{-17}$	0.091	H5	$6.64\cdot10^{5}$	$7.61\cdot10^{-16}$	1.374
O6	$-1.86\cdot10^{7}$	$4.58\cdot10^{-15}$	8.273	C6	$-2.42\cdot10^{6}$	$1.75\cdot10^{-16}$	0.316	H6	$-1.58\cdot10^{6}$	$1.81\cdot10^{-15}$	3.275
O7	$-4.83\cdot10^{6}$	$1.19\cdot10^{-15}$	2.143	C7	$-5.51\cdot10^{6}$	$3.99\cdot10^{-16}$	0.721	H7	$-1.42\cdot10^{6}$	$1.63\cdot10^{-15}$	2.939
O8	$9.28\cdot10^{6}$	$2.28\cdot10^{-15}$	4.118	C8	$1.45\cdot10^{6}$	$1.05\cdot10^{-16}$	0.189	H8	$-2.66\cdot10^{6}$	$3.04\cdot10^{-15}$	5.501
O9	$-1.52\cdot10^{7}$	$3.73\cdot10^{-15}$	6.737	C9	$-8.77\cdot10^{5}$	$6.35\cdot10^{-17}$	0.115	H9	$-5.30\cdot10^{5}$	$6.07\cdot10^{-16}$	1.097
O10	$-1.45\cdot10^{6}$	$3.57\cdot10^{-16}$	0.645	C10	$2.92\cdot10^{5}$	$2.11\cdot10^{-17}$	0.038	H10	$2.18\cdot10^{5}$	$2.49\cdot10^{-16}$	0.451
O11	$-1.63\cdot10^{6}$	$4.01\cdot10^{-16}$	0.724	C11	$3.42\cdot10^{6}$	$2.48\cdot10^{-16}$	0.448	H11	$1.73\cdot10^{6}$	$1.99\cdot10^{-15}$	3.590
N1	$2.72\cdot10^{6}$	$4.34\cdot10^{-17}$	0.078	C12	$2.71\cdot10^{6}$	$1.96\cdot10^{-16}$	0.355	H12	$8.70\cdot10^{5}$	$9.96\cdot10^{-16}$	1.801
N2	$2.71\cdot10^{6}$	$4.32\cdot10^{-17}$	0.078	C13	$1.26\cdot10^{6}$	$9.14\cdot10^{-17}$	0.165	H13	$-1.08\cdot10^{5}$	$1.24\cdot10^{-16}$	0.224
N3	$6.76\cdot10^{5}$	$1.08\cdot10^{-17}$	0.019	C14	$8.21\cdot10^{3}$	$5.94\cdot10^{-19}$	0.001	H14	$-8.14\cdot10^{5}$	$9.32\cdot10^{-16}$	1.684
N4	$-4.74\cdot10^{6}$	$7.55\cdot10^{-17}$	0.136	C15	$-2.34\cdot10^{5}$	$1.69\cdot10^{-17}$	0.031	H15	$-6.01\cdot10^{6}$	$6.88\cdot10^{-15}$	12.44
N5	$-3.89\cdot10^{6}$	$6.20\cdot10^{-17}$	0.112	C16	$-3.93\cdot10^{6}$	$2.85\cdot10^{-16}$	0.515	H16	$-2.62\cdot10^{6}$	$3.00\cdot10^{-15}$	5.418
N6	$3.99\cdot10^{6}$	$6.36\cdot10^{-17}$	0.115	C17	$1.84\cdot10^{6}$	$1.33\cdot10^{-16}$	0.240	H17	$-1.46\cdot10^{6}$	$1.67\cdot10^{-15}$	3.017
				C18	$1.04\cdot10^{6}$	$7.52\cdot10^{-17}$	0.136	H18	$1.34\cdot10^{6}$	$1.54\cdot10^{-15}$	2.782
				C19	$9.48\cdot10^{5}$	$6.87\cdot10^{-17}$	0.124	H19	$6.84\cdot10^{5}$	$7.83\cdot10^{-16}$	1.415
								H20	$6.43\cdot10^{5}$	$7.36\cdot10^{-16}$	1.330
								H21	$1.80\cdot10^{6}$	$2.06\cdot10^{-15}$	3.730
								H22	$7.33\cdot10^{6}$	$8.39\cdot10^{-15}$	15.16
								H23	$7.68\cdot10^{5}$	$8.79\cdot10^{-16}$	1.589
								H24	$1.69\cdot10^{6}$	$1.93\cdot10^{-15}$	3.494

TABLE XVIII: Values of $\beta^2\sigma^{(1p)}$, expressed in ppm au, of the rotating electric polarization, expressed in $\frac{C}{m^2}$, and of the induced voltage, expressed in μV, for non-lanthanide atoms of complex **6**. 1 ppm au of $\beta^2\sigma^{(1p)} = 1.94469097 \cdot 10^{-18}\frac{m}{V}$.

Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|
O1	$-1.22\cdot10^{6}$	$3.00\cdot10^{-16}$	0.542	C1	$-2.60\cdot10^{5}$	$1.89\cdot10^{-17}$	0.034	H1	$-3.91\cdot10^{5}$	$4.47\cdot10^{-16}$	0.808
O2	$1.24\cdot10^{7}$	$3.04\cdot10^{-15}$	5.500	C2	$-8.26\cdot10^{4}$	$5.98\cdot10^{-18}$	0.011	H2	$2.45\cdot10^{4}$	$2.81\cdot10^{-17}$	0.051
O3	$6.69\cdot10^{6}$	$1.64\cdot10^{-15}$	2.968	C3	$-3.42\cdot10^{5}$	$2.47\cdot10^{-17}$	0.045	H3	$-1.08\cdot10^{6}$	$1.23\cdot10^{-15}$	2.230
O4	$-9.17\cdot10^{5}$	$2.25\cdot10^{-16}$	0.407	C4	$-1.07\cdot10^{5}$	$7.78\cdot10^{-18}$	0.014	H4	$2.00\cdot10^{5}$	$2.30\cdot10^{-16}$	0.415
O5	$-1.12\cdot10^{7}$	$2.76\cdot10^{-15}$	4.987	C5	$1.67\cdot10^{5}$	$1.21\cdot10^{-17}$	0.022	H5	$-2.65\cdot10^{4}$	$3.04\cdot10^{-17}$	0.055
O6	$-8.51\cdot10^{6}$	$2.09\cdot10^{-15}$	3.775	C6	$-4.11\cdot10^{5}$	$2.97\cdot10^{-17}$	0.054	H6	$-2.07\cdot10^{4}$	$2.37\cdot10^{-17}$	0.043
N1	$2.66\cdot10^{5}$	$4.24\cdot10^{-18}$	0.008	C7	$-1.04\cdot10^{5}$	$7.51\cdot10^{-18}$	0.014	H7	$-5.08\cdot10^{4}$	$5.81\cdot10^{-17}$	0.105
N2	$1.57\cdot10^{6}$	$2.51\cdot10^{-17}$	0.045	C8	$-8.26\cdot10^{4}$	$5.98\cdot10^{-18}$	0.011	H8	$7.81\cdot10^{5}$	$8.94\cdot10^{-16}$	1.615
N3	$-5.43\cdot10^{4}$	$8.67\cdot10^{-19}$	0.002	C9	$3.82\cdot10^{4}$	$2.77\cdot10^{-18}$	0.005	H9	$5.08\cdot10^{4}$	$5.81\cdot10^{-17}$	0.105
N4	$-5.89\cdot10^{4}$	$9.39\cdot10^{-19}$	0.002	C10	$1.60\cdot10^{6}$	$1.16\cdot10^{-16}$	0.209	H10	$6.67\cdot10^{5}$	$7.64\cdot10^{-16}$	1.381
F1	$-9.03\cdot10^{5}$	$9.16\cdot10^{-16}$	1.656	C11	$-1.04\cdot10^{5}$	$7.56\cdot10^{-18}$	0.014	H11	$-3.73\cdot10^{4}$	$4.27\cdot10^{-17}$	0.077
F2	$-1.01\cdot10^{6}$	$1.02\cdot10^{-15}$	1.844	C12	$6.07\cdot10^{4}$	$4.40\cdot10^{-18}$	0.008	H12	$2.26\cdot10^{6}$	$2.58\cdot10^{-15}$	4.670
F3	$-1.12\cdot10^{6}$	$1.14\cdot10^{-15}$	2.061	C13	$-2.02\cdot10^{6}$	$1.46\cdot10^{-16}$	0.264	H13	$-7.89\cdot10^{4}$	$9.03\cdot10^{-17}$	0.163
F4	$-2.92\cdot10^{5}$	$2.96\cdot10^{-16}$	0.535	C14	$2.17\cdot10^{5}$	$1.57\cdot10^{-17}$	0.028	H14	$5.68\cdot10^{4}$	$6.50\cdot10^{-17}$	0.117
F5	$-7.59\cdot10^{4}$	$7.70\cdot10^{-17}$	0.139	C15	$5.99\cdot10^{5}$	$4.34\cdot10^{-17}$	0.078	H15	$9.46\cdot10^{5}$	$1.08\cdot10^{-15}$	1.958
F6	$7.84\cdot10^{4}$	$7.95\cdot10^{-17}$	0.144	C16	$-3.12\cdot10^{4}$	$2.26\cdot10^{-18}$	0.004	H16	$2.77\cdot10^{4}$	$3.17\cdot10^{-17}$	0.057
F7	$-6.26\cdot10^{5}$	$6.35\cdot10^{-16}$	1.148	C17	$6.50\cdot10^{4}$	$4.70\cdot10^{-18}$	0.009	H17	$2.55\cdot10^{5}$	$2.92\cdot10^{-16}$	0.527
F8	$-1.23\cdot10^{6}$	$1.25\cdot10^{-15}$	2.254	C18	$-2.76\cdot10^{4}$	$1.99\cdot10^{-18}$	0.004	H18	$1.06\cdot10^{6}$	$1.21\cdot10^{-15}$	2.183
F9	$-7.37\cdot10^{5}$	$7.47\cdot10^{-16}$	1.351	C19	$-1.42\cdot10^{4}$	$1.02\cdot10^{-18}$	0.002	H19	$2.70\cdot10^{5}$	$3.09\cdot10^{-16}$	0.558
				C20	$-4.77\cdot10^{4}$	$3.45\cdot10^{-18}$	0.006	H20	$3.70\cdot10^{5}$	$4.24\cdot10^{-16}$	0.767
				C21	$4.08\cdot10^{6}$	$2.95\cdot10^{-16}$	0.533	H21	$5.85\cdot10^{4}$	$6.70\cdot10^{-17}$	0.121
				C22	$1.54\cdot10^{6}$	$1.11\cdot10^{-16}$	0.201	H22	$1.59\cdot10^{6}$	$1.82\cdot10^{-15}$	3.285
				C23	$6.04\cdot10^{4}$	$4.37\cdot10^{-18}$	0.008	H23	$-1.78\cdot10^{4}$	$2.03\cdot10^{-17}$	0.037
				C24	$7.91\cdot10^{5}$	$5.73\cdot10^{-17}$	0.104	H24	$-1.91\cdot10^{4}$	$2.18\cdot10^{-17}$	0.039
				C25	$-5.66\cdot10^{4}$	$4.10\cdot10^{-18}$	0.007	H25	$3.32\cdot10^{5}$	$3.80\cdot10^{-16}$	0.687
				C26	$-4.41\cdot10^{6}$	$3.19\cdot10^{-16}$	0.577	H26	$8.84\cdot10^{4}$	$1.01\cdot10^{-16}$	0.183
				C27	$-4.57\cdot10^{4}$	$3.31\cdot10^{-18}$	0.006	H27	$2.01\cdot10^{5}$	$2.30\cdot10^{-16}$	0.416
				C28	$-1.60\cdot10^{6}$	$1.16\cdot10^{-16}$	0.210	H28	$4.55\cdot10^{5}$	$5.22\cdot10^{-16}$	0.943
				C29	$1.50\cdot10^{6}$	$1.09\cdot10^{-16}$	0.197	C39	$4.24\cdot10^{5}$	$3.07\cdot10^{-17}$	0.055
				C30	$1.38\cdot10^{5}$	$1.00\cdot10^{-17}$	0.018	C40	$-2.82\cdot10^{5}$	$2.04\cdot10^{-17}$	0.037
				C31	$9.90\cdot10^{4}$	$7.16\cdot10^{-18}$	0.013	C41	$7.98\cdot10^{4}$	$5.78\cdot10^{-18}$	0.010
				C32	$6.50\cdot10^{5}$	$4.71\cdot10^{-17}$	0.085	C42	$9.85\cdot10^{5}$	$7.13\cdot10^{-17}$	0.129
				C33	$-8.82\cdot10^{5}$	$6.39\cdot10^{-17}$	0.115	C43	$-2.28\cdot10^{4}$	$1.65\cdot10^{-18}$	0.003
				C34	$3.63\cdot10^{4}$	$2.63\cdot10^{-18}$	0.005	C44	$-2.40\cdot10^{4}$	$1.73\cdot10^{-18}$	0.003
				C35	$3.77\cdot10^{5}$	$2.73\cdot10^{-17}$	0.049	C45	$4.57\cdot10^{5}$	$3.31\cdot10^{-17}$	0.060
				C36	$1.05\cdot10^{6}$	$7.62\cdot10^{-17}$	0.138	C46	$1.68\cdot10^{5}$	$1.22\cdot10^{-17}$	0.022
				C37	$-9.26\cdot10^{5}$	$6.70\cdot10^{-17}$	0.121	C47	$2.68\cdot10^{5}$	$1.94\cdot10^{-17}$	0.035
				C38	$4.61\cdot10^{5}$	$3.34\cdot10^{-17}$	0.060	C48	$4.98\cdot10^{5}$	$3.61\cdot10^{-17}$	0.065

TABLE XIX: Values of $\beta^2\sigma^{(1p)}$, expressed in ppm au, of the rotating electric polarization, expressed in $\frac{C}{m^2}$, and of the induced voltage, expressed in μV, for non-lanthanide atoms of complex **7**. 1 ppm au of $\beta^2\sigma^{(1p)}$ $= 1.94469097 \cdot 10^{-18}\frac{m}{V}$.

Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|
O1	$1.94\cdot10^{6}$	$4.77\cdot10^{-16}$	0.862	C1	$2.13\cdot10^{4}$	$1.54\cdot10^{-18}$	0.003	H1	$5.10\cdot10^{4}$	$5.84\cdot10^{-17}$	0.105
O2	$-8.17\cdot10^{6}$	$2.00\cdot10^{-15}$	3.622	C2	$-1.02\cdot10^{5}$	$7.41\cdot10^{-18}$	0.013	H2	$-8.25\cdot10^{4}$	$9.45\cdot10^{-17}$	0.171
O3	$-1.30\cdot10^{7}$	$3.19\cdot10^{-15}$	5.773	C3	$-3.52\cdot10^{5}$	$2.55\cdot10^{-17}$	0.046	H3	$-4.32\cdot10^{5}$	$4.95\cdot10^{-16}$	0.894
O4	$1.89\cdot10^{7}$	$4.65\cdot10^{-15}$	8.399	C4	$-5.22\cdot10^{5}$	$3.78\cdot10^{-17}$	0.068	H4	$-2.28\cdot10^{6}$	$2.61\cdot10^{-15}$	4.720
O5	$1.29\cdot10^{6}$	$3.16\cdot10^{-16}$	0.571	C5	$-2.43\cdot10^{6}$	$1.76\cdot10^{-16}$	0.318	H5	$-8.28\cdot10^{5}$	$9.48\cdot10^{-16}$	1.713
O6	$-1.08\cdot10^{7}$	$2.64\cdot10^{-15}$	4.774	C6	$-3.95\cdot10^{6}$	$2.86\cdot10^{-16}$	0.517	H6	$-6.83\cdot10^{5}$	$7.82\cdot10^{-16}$	1.413
N1	$-7.13\cdot10^{6}$	$1.14\cdot10^{-16}$	0.205	C7	$-5.64\cdot10^{6}$	$4.09\cdot10^{-16}$	0.738	H7	$-9.57\cdot10^{5}$	$1.10\cdot10^{-15}$	1.980
N2	$1.39\cdot10^{6}$	$2.21\cdot10^{-17}$	0.040	C8	$-2.09\cdot10^{6}$	$1.51\cdot10^{-16}$	0.273	H8	$5.57\cdot10^{4}$	$6.38\cdot10^{-17}$	0.115
F1	$-1.41\cdot10^{6}$	$1.43\cdot10^{-15}$	2.580	C9	$-1.14\cdot10^{6}$	$8.28\cdot10^{-17}$	0.150	H9	$-1.85\cdot10^{6}$	$2.12\cdot10^{-15}$	3.831
F2	$-5.59\cdot10^{5}$	$5.67\cdot10^{-16}$	1.024	C10	$-9.90\cdot10^{5}$	$7.17\cdot10^{-17}$	0.130	H10	$-8.74\cdot10^{4}$	$1.00\cdot10^{-16}$	0.181
F3	$-1.91\cdot10^{6}$	$1.94\cdot10^{-15}$	3.503	C11	$-1.36\cdot10^{6}$	$9.82\cdot10^{-17}$	0.177	H11	$2.39\cdot10^{2}$	$2.74\cdot10^{-19}$	<0.001
F4	$1.35\cdot10^{6}$	$1.37\cdot10^{-15}$	2.467	C12	$-2.79\cdot10^{6}$	$2.02\cdot10^{-16}$	0.365	H12	$-3.26\cdot10^{5}$	$3.73\cdot10^{-16}$	0.674
F5	$6.93\cdot10^{5}$	$7.03\cdot10^{-16}$	1.271	C13	$-4.07\cdot10^{6}$	$2.94\cdot10^{-16}$	0.532	H13	$-8.40\cdot10^{5}$	$9.62\cdot10^{-16}$	1.738
F6	$1.86\cdot10^{6}$	$1.89\cdot10^{-15}$	3.408	C14	$5.02\cdot10^{5}$	$3.64\cdot10^{-17}$	0.066	H14	$-8.38\cdot10^{5}$	$9.60\cdot10^{-16}$	1.735
F7	$-1.89\cdot10^{6}$	$1.92\cdot10^{-15}$	3.469	C15	$5.22\cdot10^{6}$	$3.78\cdot10^{-16}$	0.683	H15	$-1.15\cdot10^{6}$	$1.32\cdot10^{-15}$	2.379
F8	$-4.38\cdot10^{5}$	$4.45\cdot10^{-16}$	0.803	C16	$2.00\cdot10^{6}$	$1.45\cdot10^{-16}$	0.262	H16	$-2.94\cdot10^{6}$	$3.36\cdot10^{-15}$	6.078
F9	$-1.20\cdot10^{6}$	$1.21\cdot10^{-15}$	2.192	C17	$-2.32\cdot10^{5}$	$1.68\cdot10^{-17}$	0.030	H17	$7.56\cdot10^{4}$	$8.66\cdot10^{-17}$	0.157
S1	$1.64\cdot10^{5}$	$5.56\cdot10^{-18}$	0.010	C18	$2.01\cdot10^{3}$	$1.45\cdot10^{-19}$	<0.001	H18	$-1.38\cdot10^{5}$	$1.58\cdot10^{-16}$	0.286
S2	$-2.42\cdot10^{6}$	$8.18\cdot10^{-17}$	0.148	C19	$1.55\cdot10^{4}$	$1.12\cdot10^{-18}$	0.002	H19	$-6.71\cdot10^{5}$	$7.69\cdot10^{-16}$	1.389
S3	$2.89\cdot10^{5}$	$9.78\cdot10^{-18}$	0.018	C20	$-2.21\cdot10^{5}$	$1.60\cdot10^{-17}$	0.029	H20	$-2.11\cdot10^{6}$	$2.42\cdot10^{-15}$	4.369
				C21	$-5.94\cdot10^{5}$	$4.30\cdot10^{-17}$	0.078				
				C22	$-2.66\cdot10^{6}$	$1.92\cdot10^{-16}$	0.348				
				C23	$-1.41\cdot10^{6}$	$1.02\cdot10^{-16}$	0.184				
				C24	$-1.29\cdot10^{6}$	$9.37\cdot10^{-17}$	0.169				
				C25	$-1.75\cdot10^{6}$	$1.26\cdot10^{-16}$	0.228				
				C26	$-3.81\cdot10^{6}$	$2.76\cdot10^{-16}$	0.498				
				C27	$2.65\cdot10^{4}$	$1.92\cdot10^{-18}$	0.003				
				C28	$-1.74\cdot10^{5}$	$1.26\cdot10^{-17}$	0.023				
				C29	$-5.58\cdot10^{5}$	$4.04\cdot10^{-17}$	0.073				
				C30	$-6.39\cdot10^{5}$	$4.63\cdot10^{-17}$	0.084				
				C31	$-2.65\cdot10^{6}$	$1.91\cdot10^{-16}$	0.346				
				C32	$-3.73\cdot10^{6}$	$2.70\cdot10^{-16}$	0.488				
				C33	$-5.36\cdot10^{6}$	$3.88\cdot10^{-16}$	0.701				
				C34	$-1.90\cdot10^{6}$	$1.38\cdot10^{-16}$	0.249				

TABLE XX: Values of $\beta^2\sigma^{(1p)}$, expressed in ppm au, of the rotating electric polarization, expressed in $\frac{C}{m^2}$, and of the induced voltage, expressed in μV, for non-lanthanide atoms of complex **8**. 1 ppm au of $\beta^2\sigma^{(1p)}$ $= 1.94469097 \cdot 10^{-18}\frac{m}{V}$.

Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|
O1	$3.52\cdot10^{6}$	$8.64\cdot10^{-16}$	1.561	C1	$2.55\cdot10^{5}$	$1.85\cdot10^{-17}$	0.033	H1	$2.06\cdot10^{5}$	$2.36\cdot10^{-16}$	0.427
O2	$-1.30\cdot10^{7}$	$3.20\cdot10^{-15}$	5.780	C2	$1.34\cdot10^{5}$	$9.73\cdot10^{-18}$	0.018	H2	$8.52\cdot10^{4}$	$9.75\cdot10^{-17}$	0.176
O3	$6.14\cdot10^{6}$	$1.51\cdot10^{-15}$	2.723	C3	$4.00\cdot10^{4}$	$2.90\cdot10^{-18}$	0.005	H3	$-9.11\cdot10^{4}$	$1.04\cdot10^{-16}$	0.189
O4	$-9.23\cdot10^{6}$	$2.27\cdot10^{-15}$	4.094	C4	$3.84\cdot10^{5}$	$2.78\cdot10^{-17}$	0.050	H4	$-7.40\cdot10^{5}$	$8.47\cdot10^{-16}$	1.530
O5	$-1.42\cdot10^{7}$	$3.48\cdot10^{-15}$	6.284	C5	$5.01\cdot10^{2}$	$3.63\cdot10^{-20}$	<0.001	H5	$3.30\cdot10^{5}$	$3.78\cdot10^{-16}$	0.683
O6	$2.31\cdot10^{7}$	$5.67\cdot10^{-15}$	10.25	C6	$-1.60\cdot10^{6}$	$1.16\cdot10^{-16}$	0.209	H6	$5.79\cdot10^{4}$	$6.63\cdot10^{-17}$	0.120
N1	$-6.12\cdot10^{6}$	$9.76\cdot10^{-17}$	0.176	C7	$-4.22\cdot10^{6}$	$3.06\cdot10^{-16}$	0.552	H7	$-3.58\cdot10^{5}$	$4.10\cdot10^{-16}$	0.740
N2	$1.22\cdot10^{6}$	$1.94\cdot10^{-17}$	0.035	C8	$-1.68\cdot10^{6}$	$1.21\cdot10^{-16}$	0.219	H8	$-2.02\cdot10^{6}$	$2.31\cdot10^{-15}$	4.175
F1	$-4.69\cdot10^{5}$	$4.75\cdot10^{-16}$	0.859	C9	$3.85\cdot10^{5}$	$2.79\cdot10^{-17}$	0.050	H9	$-7.18\cdot10^{5}$	$8.22\cdot10^{-16}$	1.486
F2	$-8.54\cdot10^{5}$	$8.66\cdot10^{-16}$	1.566	C10	$1.32\cdot10^{5}$	$9.53\cdot10^{-18}$	0.017	H10	$-3.98\cdot10^{5}$	$4.56\cdot10^{-16}$	0.823
F3	$-2.02\cdot10^{6}$	$2.05\cdot10^{-15}$	3.711	C11	$-9.30\cdot10^{4}$	$6.73\cdot10^{-18}$	0.012	H11	$-7.99\cdot10^{4}$	$9.15\cdot10^{-17}$	0.165
F4	$-1.22\cdot10^{6}$	$1.24\cdot10^{-15}$	2.244	C12	$2.59\cdot10^{5}$	$1.87\cdot10^{-17}$	0.034	H12	$1.74\cdot10^{6}$	$2.00\cdot10^{-15}$	3.605
F5	$-3.24\cdot10^{5}$	$3.29\cdot10^{-16}$	0.595	C13	$-1.10\cdot10^{6}$	$7.98\cdot10^{-17}$	0.144	H13	$-1.72\cdot10^{6}$	$1.97\cdot10^{-15}$	3.554
F6	$-1.62\cdot10^{6}$	$1.65\cdot10^{-15}$	2.976	C14	$-3.34\cdot10^{6}$	$2.42\cdot10^{-16}$	0.437	H14	$-1.60\cdot10^{5}$	$1.83\cdot10^{-16}$	0.330
F7	$1.76\cdot10^{6}$	$1.79\cdot10^{-15}$	3.231	C15	$-5.08\cdot10^{6}$	$3.68\cdot10^{-16}$	0.664	H15	$1.12\cdot10^{5}$	$1.28\cdot10^{-16}$	0.231
F8	$1.59\cdot10^{6}$	$1.61\cdot10^{-15}$	2.918	C16	$-1.76\cdot10^{6}$	$1.27\cdot10^{-16}$	0.230	H16	$1.53\cdot10^{5}$	$1.75\cdot10^{-16}$	0.316
F9	$2.68\cdot10^{6}$	$2.72\cdot10^{-15}$	4.910	C17	$-9.15\cdot10^{5}$	$6.63\cdot10^{-17}$	0.120	H17	$5.28\cdot10^{4}$	$6.05\cdot10^{-17}$	0.109
S1	$6.55\cdot10^{5}$	$2.21\cdot10^{-17}$	0.040	C18	$-5.93\cdot10^{5}$	$4.29\cdot10^{-17}$	0.078	H18	$-2.65\cdot10^{5}$	$3.03\cdot10^{-16}$	0.548
S2	$1.10\cdot10^{6}$	$3.71\cdot10^{-17}$	0.067	C19	$-4.28\cdot10^{5}$	$3.10\cdot10^{-17}$	0.056	H19	$-8.44\cdot10^{5}$	$9.66\cdot10^{-16}$	1.746
S3	$-2.16\cdot10^{6}$	$7.29\cdot10^{-17}$	0.132	C20	$-1.44\cdot10^{6}$	$1.04\cdot10^{-16}$	0.189	H20	$-4.51\cdot10^{6}$	$5.17\cdot10^{-15}$	9.338
				C21	$-1.03\cdot10^{6}$	$7.44\cdot10^{-17}$	0.134				
				C22	$3.31\cdot10^{6}$	$2.40\cdot10^{-16}$	0.434				
				C23	$8.18\cdot10^{6}$	$5.92\cdot10^{-16}$	1.070				
				C24	$3.14\cdot10^{6}$	$2.27\cdot10^{-16}$	0.410				
				C25	$-4.92\cdot10^{5}$	$3.56\cdot10^{-17}$	0.064				
				C26	$-4.79\cdot10^{4}$	$3.47\cdot10^{-18}$	0.006				
				C27	$1.79\cdot10^{5}$	$1.29\cdot10^{-17}$	0.023				
				C28	$3.89\cdot10^{5}$	$2.81\cdot10^{-17}$	0.051				
				C29	$8.25\cdot10^{5}$	$5.97\cdot10^{-17}$	0.108				
				C30	$2.11\cdot10^{5}$	$1.53\cdot10^{-17}$	0.028				
				C31	$9.73\cdot10^{4}$	$7.04\cdot10^{-18}$	0.013				
				C32	$-1.83\cdot10^{5}$	$1.32\cdot10^{-17}$	0.024				
				C33	$-3.97\cdot10^{5}$	$2.87\cdot10^{-17}$	0.052				
				C34	$-4.61\cdot10^{5}$	$3.34\cdot10^{-17}$	0.060				
				C35	$-1.11\cdot10^{6}$	$8.04\cdot10^{-17}$	0.145				
				C36	$-3.63\cdot10^{6}$	$2.63\cdot10^{-16}$	0.475				

TABLE XXI: Values of $\beta^2\sigma^{(1p)}$, expressed in ppm au, of the rotating electric polarization, expressed in $\frac{C}{m^2}$, and of the induced voltage, expressed in μV, for non-lanthanide atoms of complex **9**. 1 ppm au of $\beta^2\sigma^{(1p)}$ = $1.94469097\cdot10^{-18}\frac{m}{V}$.

Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|	Atom	$\beta^2\sigma^{(1p)}$	|P|	|V|
O1	$-4.71\cdot10^{6}$	$1.16\cdot10^{-15}$	2.087	C1	$2.62\cdot10^{6}$	$1.90\cdot10^{-16}$	0.343	H1	$4.62\cdot10^{5}$	$5.29\cdot10^{-16}$	0.956
O2	$6.28\cdot10^{6}$	$1.54\cdot10^{-15}$	2.787	C2	$5.93\cdot10^{5}$	$4.29\cdot10^{-17}$	0.078	H2	$1.37\cdot10^{5}$	$1.57\cdot10^{-16}$	0.284
O3	$-7.83\cdot10^{6}$	$1.92\cdot10^{-15}$	3.475	C3	$4.05\cdot10^{5}$	$2.93\cdot10^{-17}$	0.053	H3	$-5.27\cdot10^{4}$	$6.03\cdot10^{-17}$	0.109
O4	$1.60\cdot10^{7}$	$3.92\cdot10^{-15}$	7.088	C4	$1.58\cdot10^{5}$	$1.14\cdot10^{-17}$	0.021	H4	$2.61\cdot10^{6}$	$2.99\cdot10^{-15}$	5.402
O5	$1.87\cdot10^{7}$	$4.59\cdot10^{-15}$	8.300	C5	$-6.45\cdot10^{3}$	$4.67\cdot10^{-19}$	0.001	H5	$7.83\cdot10^{5}$	$8.97\cdot10^{-16}$	1.621
O6	$-2.63\cdot10^{7}$	$6.46\cdot10^{-15}$	11.68	C6	$4.25\cdot10^{6}$	$3.08\cdot10^{-16}$	0.556	H6	$1.91\cdot10^{5}$	$2.19\cdot10^{-16}$	0.396
N1	$-3.66\cdot10^{5}$	$5.84\cdot10^{-18}$	0.011	C7	$5.03\cdot10^{6}$	$3.65\cdot10^{-16}$	0.659	H7	$-1.73\cdot10^{5}$	$1.98\cdot10^{-16}$	0.358
N2	$7.47\cdot10^{6}$	$1.19\cdot10^{-16}$	0.215	C8	$1.65\cdot10^{6}$	$1.20\cdot10^{-16}$	0.216	H8	$2.83\cdot10^{6}$	$3.24\cdot10^{-15}$	5.850
F1	$1.31\cdot10^{6}$	$1.33\cdot10^{-15}$	2.408	C9	$2.33\cdot10^{6}$	$1.68\cdot10^{-16}$	0.304	H9	$-1.85\cdot10^{5}$	$2.12\cdot10^{-16}$	0.383
F2	$4.52\cdot10^{5}$	$4.59\cdot10^{-16}$	0.829	C10	$6.12\cdot10^{5}$	$4.43\cdot10^{-17}$	0.080	H10	$3.20\cdot10^{5}$	$3.67\cdot10^{-16}$	0.662
F3	$1.15\cdot10^{6}$	$1.16\cdot10^{-15}$	2.105	C11	$6.54\cdot10^{5}$	$4.73\cdot10^{-17}$	0.086	H11	$7.64\cdot10^{5}$	$8.75\cdot10^{-16}$	1.581
F4	$-2.16\cdot10^{6}$	$2.19\cdot10^{-15}$	3.956	C12	$2.11\cdot10^{5}$	$1.52\cdot10^{-17}$	0.028	H12	$-2.20\cdot10^{6}$	$2.52\cdot10^{-15}$	4.559
F5	$-2.09\cdot10^{6}$	$2.12\cdot10^{-15}$	3.826	C13	$-1.39\cdot10^{5}$	$1.01\cdot10^{-17}$	0.018	H13	$2.61\cdot10^{6}$	$2.99\cdot10^{-15}$	5.402
F6	$-3.05\cdot10^{6}$	$3.10\cdot10^{-15}$	5.593	C14	$4.80\cdot10^{6}$	$3.47\cdot10^{-16}$	0.628	H14	$1.30\cdot10^{5}$	$1.49\cdot10^{-16}$	0.270
F7	$1.26\cdot10^{6}$	$1.28\cdot10^{-15}$	2.317	C15	$7.14\cdot10^{6}$	$5.17\cdot10^{-16}$	0.934	H15	$6.67\cdot10^{3}$	$7.64\cdot10^{-18}$	0.014
F8	$1.53\cdot10^{6}$	$1.55\cdot10^{-15}$	2.806	C16	$2.65\cdot10^{6}$	$1.92\cdot10^{-16}$	0.347	H16	$2.81\cdot10^{5}$	$3.22\cdot10^{-16}$	0.582
F9	$2.41\cdot10^{6}$	$2.45\cdot10^{-15}$	4.424	C17	$1.63\cdot10^{6}$	$1.18\cdot10^{-16}$	0.213	H17	$7.51\cdot10^{5}$	$8.60\cdot10^{-16}$	1.554
S1	$-1.87\cdot10^{5}$	$6.33\cdot10^{-18}$	0.011	C18	$1.72\cdot10^{6}$	$1.25\cdot10^{-16}$	0.225	H18	$3.67\cdot10^{6}$	$4.20\cdot10^{-15}$	7.587
S2	$-7.39\cdot10^{5}$	$2.50\cdot10^{-17}$	0.045	C19	$3.71\cdot10^{5}$	$2.69\cdot10^{-17}$	0.049	H19	$2.14\cdot10^{5}$	$2.45\cdot10^{-16}$	0.443
S3	$2.54\cdot10^{6}$	$8.59\cdot10^{-17}$	0.155	C20	$5.40\cdot10^{5}$	$3.91\cdot10^{-17}$	0.071	H20	$6.37\cdot10^{5}$	$7.29\cdot10^{-16}$	1.318
				C21	$9.61\cdot10^{5}$	$6.96\cdot10^{-17}$	0.126	H21	$3.24\cdot10^{5}$	$3.71\cdot10^{-16}$	0.670
				C22	$-3.84\cdot10^{6}$	$2.78\cdot10^{-16}$	0.502	H22	$7.57\cdot10^{5}$	$8.66\cdot10^{-16}$	1.566
				C23	$-9.29\cdot10^{6}$	$6.72\cdot10^{-16}$	1.215	H23	$1.01\cdot10^{5}$	$1.16\cdot10^{-16}$	0.210
				C24	$-3.75\cdot10^{6}$	$2.71\cdot10^{-16}$	0.490	H24	$-1.37\cdot10^{5}$	$1.56\cdot10^{-16}$	0.283
				C25	$7.73\cdot10^{5}$	$5.60\cdot10^{-17}$	0.101	H25	$1.41\cdot10^{6}$	$1.61\cdot10^{-15}$	2.910
				C26	$7.99\cdot10^{4}$	$5.78\cdot10^{-18}$	0.010	H26	$1.78\cdot10^{6}$	$2.04\cdot10^{-15}$	3.691
				C27	$8.87\cdot10^{3}$	$6.42\cdot10^{-19}$	0.001	H27	$9.19\cdot10^{5}$	$1.05\cdot10^{-15}$	1.902
				C28	$2.15\cdot10^{5}$	$1.56\cdot10^{-17}$	0.028	H28	$4.47\cdot10^{5}$	$5.12\cdot10^{-16}$	0.926
				C29	$6.83\cdot10^{5}$	$4.94\cdot10^{-17}$	0.089	H29	$3.95\cdot10^{5}$	$4.52\cdot10^{-16}$	0.817
				C30	$2.26\cdot10^{6}$	$1.64\cdot10^{-16}$	0.296	H30	$6.59\cdot10^{5}$	$7.55\cdot10^{-16}$	1.364
				C31	$1.19\cdot10^{6}$	$8.60\cdot10^{-17}$	0.155				
				C32	$9.88\cdot10^{5}$	$7.16\cdot10^{-17}$	0.129	C37	$8.30\cdot10^{5}$	$6.01\cdot10^{-17}$	0.109
				C33	$1.55\cdot10^{6}$	$1.12\cdot10^{-16}$	0.203	C38	$8.66\cdot10^{5}$	$6.27\cdot10^{-17}$	0.113
				C34	$3.94\cdot10^{6}$	$2.85\cdot10^{-16}$	0.516	C39	$1.68\cdot10^{5}$	$1.22\cdot10^{-17}$	0.022
				C35	$5.25\cdot10^{5}$	$3.80\cdot10^{-17}$	0.069	C40	$1.29\cdot10^{6}$	$9.37\cdot10^{-17}$	0.169
				C36	$4.60\cdot10^{5}$	$3.33\cdot10^{-17}$	0.060	C41	$5.60\cdot10^{5}$	$4.05\cdot10^{-17}$	0.073

TABLE XXII: Values of $\beta^2\sigma^{(1p)}$, expressed in ppm au, of the rotating electric polarization, expressed in $\frac{C}{m^2}$, and of the induced voltage, expressed in μV, for non-lanthanide atoms of complex **10**. 1 ppm au of $\beta^2\sigma^{(1p)}$ = $1.94469097\cdot10^{-18}\frac{m}{V}$.

* asoncini@unimelb.edu.au

[1] F. Aquilante, L. De Vico, N. Ferré, G. Ghigo, P.-Å. Malmqvist, P. Neogrády, T. B. Pedersen, M. Pitonak, M. Reiher, B. O. Roos, L. Serrano-Andrés, M. Urban, V. Veryazov, and R. Lindh, J. Comput. Chem. **31**, 224 (2010).

[2] N. F. Chilton, D. Collison, E. J. L. McInnes, R. E. P. Winpenny, and A. Soncini, Nat. Commun. **4**, 2551 (2013).

[3] L. Ungur and L. F. Chibotaru, in *Lanthanides and Actinides in Molecular Magnetism*, edited by R. A. Layfield and M. Murugesu (Wiley–VCH, 2015) pp. 153–184.

[4] A. D. Buckingham and P. Fischer, Chem. Phys. **324**, 111 (2006).

[5] S. Pelloni, F. Faglioni, and P. Lazzeretti, Rend. Fis. Acc. Lincei **24**, 283 (2013).

[6] A. D. Buckingham, P. Lazzeretti, and S. Pelloni, Mol. Phys. **113**, 1780 (2015).